

Checklist of the aphyllophoraceous fungi (*Agaricomycetes*) of the Brazilian Amazonia

ALLYNE CHRISTINA GOMES-SILVA¹ & TATIANA BAPTISTA GIBERTONI¹

allynefungi@hotmail.com tbgibertoni@hotmail.com

Universidade Federal de Pernambuco, Departamento de Micologia

Av. Nelson Chaves s/n, CEP 50760-420, Recife, PE, Brazil

Abstract — A literature-based checklist of the aphyllophoraceous fungi reported from the Brazilian Amazonia was compiled. Two hundred and sixteen species, 90 genera, 22 families, and 9 orders (*Agaricales*, *Auriculariales*, *Cantharellales*, *Corticiales*, *Gloeophyllales*, *Hymenochaetales*, *Polyporales*, *Russulales* and *Trechisporales*) have been reported from the area.

Key words — macrofungi, neotropics

Introduction

The aphyllophoraceous fungi are currently spread throughout many orders of *Agaricomycetes* (Hibbett et al. 2007) and comprise species that function as major decomposers of plant organic matter (Alexopoulos et al. 1996).

The Amazonian Forest (00°44'–06°24'S / 58°05'–68°01'W) covers an area of 7×10^6 km² in nine South American countries. Around 63% of the forest is located in nine Brazilian States (Acre, Amazonas, Amapá, Pará, Rondônia, Roraima, Tocantins, west of Maranhão, and north of Mato Grosso) (Fig. 1). The Amazonian forest consists of a mosaic of different habitats, such as open ombrophilous, stationally semi-deciduous, mountain, “terra firme,” “várzea” and “igapo” forests, and “campinaranas” (Amazonian savannahs). Six months of dry season and six months of rainy season can be observed (Museu Paraense Emílio Goeldi 2007).

Even with the high biodiversity of Amazonia and the well-documented importance of aphyllophoraceous fungi to all arboreous ecosystems, few studies have been undertaken in the Brazilian Amazonia on this group of fungi (Bononi 1981, 1992, Capelari & Maziero 1988, Gomes-Silva et al. 2008, Jesus 1996, Martin-Júnior et al. 2008, Sotão et al. 1997, 2002, 2008). This work aims to contribute to the knowledge about the diversity of the aphyllophoraceous fungi in this highly diverse biome by compiling and updating the nomenclature of published records of aphyllophoraceous fungi from the Brazilian Amazon.

Material and methods

All available papers with records of aphyllophoraceous fungi (*Agaricomycetes*) collected in the Brazilian Amazonia were examined. The reports were compiled and the nomenclature of the listed species were checked against and — when necessary— updated following the classification of Index Fungorum (<http://www.indexfungorum.org/Names/Names.asp>) and CBS (<http://www.cbs.knaw.nl/databases/>). Taxa with unresolved nomenclature according to the consulted databases and those lacking neo- or pantropical distributions were excluded. The occurrence and distribution of the species in Brazilian states were recorded. The checklist is organized in alphabetical order by families, genus, and species.

Results and discussion

Currently, 216 species of aphyllophoraceous fungi previously recorded from the Brazilian Amazonia are included in the checklist. These species are at present distributed in 90 genera, 22 families, and 9 orders (*Agaricales*, *Auriculariales*, *Cantharellales*, *Corticiales*, *Gloeophyllales*, *Hymenochaetales*, *Polyporales*, *Russulales* and *Trechisporales*). *Polyporales* has the highest number of species (146), families (7), and genera (61). Among the families, *Polyporaceae* is the most species rich (79), followed by *Hymenochaetaceae* (32), and *Meruliaceae* (29). *Botryobasidiaceae*, *Cantharellaceae*, *Corticiaceae*, *Cystostereaceae*, *Meripilaceae*, *Peniophoraceae*, *Schizophyllaceae* and *Schizoporaceae* are each represented by one species. Among the genera, *Amauroderma* (16), *Phellinus* (15), *Trametes* (14), *Hymenochaete* (8), *Ganoderma* (8), *Polystictus* (7), and *Podoscypha* (7) have the highest number of species. *Phellinus gilvus* (Schwein.) Pat. and *Earliella scabrosa* (Pers.) Gilb. & Ryvarden have been recorded in all states of the Brazilian Amazonia. The State of Pará is represented by 126 species (58%), with 53 exclusive to the State. The State of Amazonas has 86 reported species (35 exclusive) and 35 species were recorded from the State of Roraima (15 exclusive). The states of Rondônia and Acre recorded 48 and 28 species respectively, each with 7 exclusive species and the states of Mato Grosso and Amapá recorded 36 and 23 species respectively, each with 6 exclusive species. Of the excluded species, all of which should have their exsiccates reviewed, 22 were excluded due to their non neo- or pantropical distribution and 65 due to nomenclature uncertainties according to the consulted databases.

***Auriculariaceae* Fries ex Lindau**

***Auricularia auricula-judae* (Bull.) Quél.**

Distribution: Pará [Hennings (1908), Batista et al. (1966), Sotão et al. (2003), Campos et al. (2005)], as *Auricularia auricula* (L.) Underw.]

***Auricularia fuscosuccinea* (Mont.) Henn.**

Distribution: Pará (Sotão et al. 2003; Campos et al. 2005); Rondônia (Capelari & Maziero 1988)

***Auricularia mesenterica* (Dicks.) Pers.**

Distribution: Amazonas (Hennings 1904); Pará (Batista et al. 1966)

***Auricularia polytricha* (Mont.) Sacc.**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003); Mato Grosso (Fidalgo 1968); Pará (Batista et al. 1966; Sotão et al. 2002a, 2003); Rondônia (Capelari & Maziero 1988); Roraima (Jesus 1996)

***Pseudohydnum gelatinosum* (Scop.) P. Karst.**

Distribution: Pará (Bononi 1981)

***Bondarzewiaceae* Kotlába & Pouzar**

***Stecchericium isabellinum* Corner**

Distribution: Amazonas (Corner 1989)

***Stecchericium seriatum* (Lloyd) Maas Geest.**

Distribution: Amazonas (Bononi 1981)

***Wrightoporia avellanea* (Bres.) Pouzar**

Distribution: Pará (Sotão et al. 2008)

***Wrightoporia crenea* Ryvarden**

Distribution: Roraima (Ryvarden 1987)

***Botryobasidiaceae* (Parmasto) Jülich**

***Botryobasidium obtusisporum* J. Erikss.**

Distribution: Roraima (Jesus 1996)

***Cantharellaceae* J. Schrot**

***Cantharellus cibarius* Fr.**

Distribution: Amazonas (Hennings 1904)

***Clavariaceae* Chevallier**

***Clavulina connata* (Berk.) Corner**

Distribution: Amazonas (Corner 1950)

***Clavulina delicia* (Berk.) Corner**

Distribution: Amazonas (Corner 1950)

***Clavulina panurensis* (Berk.) Corner**

Distribution: Amazonas (Corner 1950)

***Clavulina sprucei* (Berk.) Corner**

Distribution: Amazonas (Corner 1950)

***Scytinopogon scaber* (Berk. & M.A. Curtis) D.A. Reid**

Distribution: Amazonas [Burt (1920), as *Stereum proliferum* (Berk.) Lloyd; Bononi (1981)]

***Corticaceae* Herter**

***Vuilleminia comedens* (Nees) Maire**

Distribution: Roraima (Jesus 1996)

***Cystostereaceae* Jülich**

***Cystidiodontia artocreas* (Berk. & M.A. Curtis ex Cooke) Hjortstam**

Distribution: Roraima (Jesus 1996)

***Fomitopsidaceae* Jülich**

***Daedalea aethalodes* (Mont.) Rajch.**

Distribution: Pará (Sotão et al. 2008)

***Fomitopsis cupreorosea* (Berk.) J. Carranza & Gilb.**

Distribution: Mato Grosso [Fidalgo (1968), as *Polyporus roseofuscus* Romell]

***Fomitella supina* (Sw.) Murrill**

Distribution: Pará (Sotão et al. 2008)

***Laetiporus persicinus* (Berk. & M.A. Curtis) Gilb.**

Distribution: Amapá [Sotão et al. (1991, 2002a, 2003), as *Amauroderma brittonii* Murrill]; Pará (Sotão et al. 2008)

***Ganodermataceae* Donk**

***Amauroderma aurantiacum* (Torrend) Gibertoni & Bernicchia**

Distribution: Rondônia (Gibertoni et al. 2008); Pará [Sotão et al. (1997, 2002b), as *Amauroderma macrosporum* J.S. Furtado]

***Amauroderma boleticeum* (Pat. & Gaillard) Torrend**

Distribution: Mato Grosso (Furtado 1981); Rondônia [Capelari & Maziero (1988), as *Amauroderma* sp. SP211529]; Pará (Furtado 1981)

***Amauroderma calcigenum* (Berk.) Torrend**

Distribution: Amazonas, Mato Grosso (Furtado 1981; Corner 1983); Pará (Sotão et al. 2008)

***Amauroderma camerarium* (Berk.) J.S. Furtado**

Distribution: Amazonas, Mato Grosso (Furtado 1981); Pará (Sotão et al. 2008)

***Amauroderma exile* (Berk.) Torrend**

Distribution: Amazonas [Furtado (1981), Corner (1983), Torrend (1920, 1922), as *Amauroderma renatum* (Berk.) Murrill; Hennings (1904), as *Polystictus parvula* (Berk.) Cooke, *P. marasmoides* (Berk.) Cooke and *P. renatus* (Berk.) Cooke]; Pará (Sotão et al. 2008)

***Amauroderma fasciculatum* (Pat.) Torrend**

Distribution: Acre (Bononi 1992)

***Amauroderma intermedium* (Bres. & Pat.) Torrend**

Distribution: Pará (Sotão et al. 2008)

***Amauroderma omphalodes* (Berk.) Torrend**

Distribution: Amazonas [Torrend (1920, 1922); Hennings (1904), as *Fomes omphalodes* (Berk.) Sacc.; Furtado 1981]; Pará (Sotão et al. 2008); Rondônia [Capelari & Maziero (1988), as *Amauroderma* sp. SP211532]

***Amauroderma partitum* (Berk.) Wakef.**

Distribution: Pará (Sotão et al. 2008)

***Amauroderma praetervisum* (Pat.) Torrend**

Distribution: Mato Grosso, Pará (Furtado 1981)

***Amauroderma subrugosum* (Bres. & Pat.) Torrend**

Distribution: Pará (Sotão et al. 1997, 2002b)

***Amauroderma schomburgkii* (Mont. & Berk.) Torrend**

Distribution: Amazonas [Furtado (1981), Corner (1983), Torrend (1920, 1922), as *Amauroderma heteromorphum* (Lév.) Torrend and *A. ocellatum* (Berk.) Lloyd ex Wakef.]; Mato Grosso (Furtado 1981); Pará (Furtado 1981; Sotão et al. 2002b, 2008); Rondônia (Capelari & Maziero 1988).

***Amauroderma sprucei* (Pat.) Torrend**

Distribution: Amazonas (Furtado 1981; Corner 1983; Torrend 1920, 1922); Pará [Furtado (1981); Sotão et al. (2002b); Sotão et al. (2008), as *Amauroderma dubiopansum* (Lloyd) Dennis]

***Amauroderma trichodematum* J.S. Furtado**

Distribution: Pará (Furtado 1981)

***Amauroderma unilaterum* (Lloyd) Ryvarden**

Distribution: Amazonas (Torrend 1920, 1922)

***Amauroderma variabile* (Berk.) Lloyd ex Wakef.**

Distribution: Amazonas (Torrend 1920, 1922)

***Ganoderma amazonense* Weir**

Distribution: Amazonas, Pará e Mato Grosso e Rondônia (Furtado 1967; Steyaert 1980)

***Ganoderma australe* (Fr.) Pat.**

Distribution: Pará [Silva & Minter (1995), Batista et al. (1966), as *Fomes australis* Fr.; Sotão et al. (1997, 2002b, 2008)]; Rondônia [Capelari & Maziero (1988), as *Ganoderma* sp. SP211485, SP211487 and *G. applanatum* (Pers.) Pat.]

***Ganoderma elegantum* Ryvarden**

Distribution: Rondônia [Capelari & Maziero (1988), as *Ganoderma lucidum* (Mont.) Pat., SP211483]

***Ganoderma multiplicatum* (Mont.) Pat.**

Distribution: Amazonas (Steyaert 1980); Pará (Matins-Júnior 2007); Rondônia [Capelari e Maziero (1988), as *Ganoderma lucidum* (Mont.) Pat., SP211484]

***Ganoderma multiplicatum* var. *vitalii* Steyaert**

Distribution: Pará (Steyaert 1962)

***Ganoderma nitidum* Murrill**

Distribution: Amazonas (Corner 1983)

***Ganoderma resinaceum* Boud.**

Distribution: Pará (Sotão et al. 2008)

***Ganoderma stipitatum* (Murrill) Murrill**

Distribution: Pará (Sotão et al. 2008); Rondônia [Capelari & Maziero (1988), as *Ganoderma* sp. SP211614, *G. lucidum* (Mont.) Pat. SP211486 and *G. subfornicatum* Murrill SP214815]

***Haddowia longipes* (Lév.) Steyaert**

Distribution: Pará (Sotão et al. 2008, as *Amauroderma renidens* (Bres.) Torrend)

***Haddowia neurospora* (J.S. Furtado) Teixeira**

Distribution: Mato Grosso [Furtado (1967), as *Ganoderma neurosporum* J.S. Furtado]

***Humphreya coffeata* (Berk.) Steyaert**

Distribution: Amazonas [Furtado (1967), as *Ganoderma coffeatum* (Berk.) J.S. Furtado]; Pará (Sotão et al. 2008)

***Gloeophyllaceae* Jülich**

***Gloeophyllum striatum* (Fr.) Murrill**

Distribution: Mato Grosso [Theissen (1911), as *Lenzites striata* (Sw.) Fr.; Fidalgo (1968)]; Pará (Sotão et al. 2002a; Campos et al. 2005)

***Gloeophyllum mexicanum* (Mont.) Ryvarden**

Distribution: Amapá (Sotão et al. 1991, 2002a)

***Mycobonia flava* (Sw.) Fr.**

Distribution: Amazonas [Hennings (1904), Theissen (1912), as *Bonia flava* (Sw.) Pat.; Bononi et al. (1981); Hjoststam & Bononi (1986)]

***Hericiaceae* Donk**

***Dentipellis fragilis* (Pers.) Donk**

Distribution: Pará (Sotão et al. 2008)

***Laxitextum bicolor* (Pers.) Lentz**

Distribution: Pará [Sotão et al. (2002a, 2003), as *Stereum bicolor* (Pers.) Pat.]

***Hydnodontaceae* Jülich**

***Hydnodon thelephorum* (Lév.) Banker**

Distribution: Amazonas (Bononi 1981)

***Trechispora farinacea* (Pers.) Liberta**

Distribution: Pará (Sotão et al. 2008)

***Hymenochaetaceae* Imazeki & Toki**

***Coltricia cinnamomea* (Pers.) Murrill**

Distribution: Amazonas [Hennings (1904), as *Polystictus oblectans* (Berk.) Cooke and *P. bulbipes* (Fr.) Fr.]; Pará (Sotão et al. 2008)

***Coltricia fragilissima* (Mont.) Ryvarden**

Distribution: Rondônia (Gomes-Silva et al. 2008)

***Coltricia hamata* (Romell) Ryvarden**

Distribution: Amazonas [Fidalgo 1968, as *Polystictus oblivionis* Lloyd]; Mato Grosso [Torrend (1926), as *Pelloporus hamatus* Romell Fidalgo (1968), as *Pelloporus hamatus* Romell]

***Fuscoporia senex* (Nees & Mont.) Ghobad-Nejhad**

Distribution: Amazonas [Sousa (1980), as *Phellinus senex* (Nees & Mont.) Imazeki]; Pará [Sotão et al. (2003), as *P. senex*)

***Hymenochaete cacao* (Berk.) Berk. & M.A. Curtis**

Distribution: Roraima (Jesus 1996)

***Hymenochaete cinnamomea* (Pers.) Bres**

Distribution: Mato Grosso (Gibertoni 2003)

***Hymenochaete damicornis* (Link) Lév.**

Distribution: Acre (Gibertoni 2003); Amazonas (Hennings 1904; Gibertoni 2003); Mato Grosso (Gibertoni 2003)

***Hymenochaete leonina* Berk. & M.A. Curtis**

Distribution: Amazonas, Rondônia (Léger 1998; Gibertoni 2003); Roraima (Gibertoni 2003)

***Hymenochaete luteobadia* (Fr.) Höhn. & Litsch**

Distribution: Amazonas (Gibertoni 2003); Mato Grosso (Fidalgo 1968; Gibertoni 2003; Pará (Léger 1998)

***Hymenochaete pinnatifida* Burt**

Distribution: Amazonas (Gibertoni 2003)

***Hymenochaete reniformis* (Fr.) Lév.**

Distribution: Acre, Mato Grosso (Gibertoni 2003)

***Hymenochaete rheicolor* (Mont.) Lév.**

Distribution: Rondônia (Gibertoni 2003)

***Inonotus luteoumbrinus* (Romell) Ryvarden**

Distribution: Acre [Bononi (1992), as *Phellinus sublinteus* (Murrill) Ryvarden]

***Inonotus tabacinus* (Mont.) G. Cunn.**

Distribution: Mato Grosso (Theissen 1911)

***Phellinus baccharidis* (Pat.) Pat.**

Distribution: Pará (Sotão et al. 2008)

***Phellinus calcitratus* (Berk. & M.A. Curtis) Ryvarden**

Distribution: Amazonas (Sousa 1980); Pará (Sotão et al. 2008)

***Phellinus dependens* (Murrill) Ryvarden**

Distribution: Amazonas (Sousa 1980)

***Phellinus extensus* (Lév.) Pat.**

Distribution: Amazonas (Sousa 1980); Pará (Sousa 1980; Sotão et al. 2008).

***Phellinus fastuosus* (Lév.) Ryvarden**

Distribution: Amazonas, Mato Grosso, Rondônia, Roraima (Sousa 1980); Pará (Sousa 1980; Sotão et al. 2003, 2008; Campos et al. 2005)

***Phellinus ferrugineovelutinus* (Henn.) Ryvarden**

Distribution: Acre [Batista et al. (1966), as *Poria ferrugineovelutina* P. Henn.]; Amazonas [Hennings (1905), as *P. ferrugineovelutina*]

***Phellinus gilvus* (Schwein.) Pat.**

Distribution: Acre (Bononi 1992); Amapá (Sotão et al. 1991, 2002a, 2003); Amazonas [Hennings (1904), Theissen (1911), as *Polyporus gilvus* (Schwein.) Fr. and *P. scruposus* Fr.; Batista et al. (1966), Silva & Minter (1995), as *P. gilvus*; Sousa (1980)]; Mato Grosso [Theissen (1911), as *P. gilvus* and *P. licnoides* (Mont.) Fr.; Fidalgo (1968); Sousa (1980)]; Pará (Sousa 1980; Sotão et al. 1997, 2002a, 2002b, 2003, 2008; Campos et al. 2005); Rondônia (Capelari & Maziero 1988); Roraima (Sousa 1980; Jesus 1996).

***Phellinus mangrovicus* (Imazeki) Imazeki**

Distribution: Pará (Campos & Cavalcanti 2000; Campos et al. 2005; Sotão et al. 2003)

***Phellinus noxius* (Corner) G. Cunn.**

Distribution: Amazonas, Mato Grosso (Sousa 1980); Roraima (Jesus 1996)

***Phellinus pseudoseneckii* (Murrill) Bondartseva & S. Herrera**

Distribution: Amazonas, Mato Grosso (Sousa 1980); Pará (Sotão et al. 2002b, 2008).

***Phellinus punctatus* (Fr.) Pilát**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003)

***Phellinus robiniae* (Murrill) A. Ames**

Distribution: Mato Grosso (Sousa 1980)

***Phellinus robustus* (Karst.) Bourd. & Galz**

Distribution: Amazonas, Mato Grosso (Sousa 1980); Pará (Sousa 1980; Sotão et al. 2002b, 2008); Rondônia (Capelari & Maziero 1988)

***Phellinus shaferi* (Murrill) Ryvarden**

Distribution: Pará (Martins-Júnior 2007)

***Phellinus undulatus* (Murrill) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Phylloporia chrysites* (Berk.) Ryvarden**

Distribution: Acre [Bononi (1992), as *Polyporus corosus* Murrill]; Pará (Sotão et al. 2008)

***Phylloporia pectinata* (Klotzsch) Ryvarden**

Distribution: Amazonas, Pará, Roraima [Sousa (1980), as *Phellinus pectinatus* (Klotzsch) Quéél.]; Mato Grosso [Theissen (1911), as *Fomes pectinatus* (Klotzsch) Gillet]

***Phylloporia spathulata* (Hook.) Ryvarden**

Distribution: Amazonas [Theissen (1911), Batista et al. (1966), Silva & Minter (1995), as *Polystictus luteonitidus* Berk.]; Rondônia [Capelari & Maziero (1988), as *Coltricia spathulata* (Hook.) Murrill]

***Lachnocladiaceae* D.A. Reid**

***Asterostroma cervicolor* (Berk. & M.A. Curtis) Massee**

Distribution: Roraima (Jesus 1996)

***Asterostroma muscicola* (Berk. & M.A. Curtis) Massee**

Distribution: Roraima (Jesus 1996)

***Lachnocladium brasiliense* (Lév.) Pat.**

Distribution: Acre (Batista et al. 1966; Silva & Minter 1995); Amazonas (Hennings 1904; Theissen 1912)

***Scytonostroma duriusculum* (Berk. & Broome) Donk**

Distribution: Roraima (Jesus 1996)

***Scytonostroma hemidichophyticum* Pouzar**

Distribution: Roraima (Jesus 1996)

***Scytonostroma odoratum* (Fr.) Donk**

Distribution: Roraima (Jesus 1996)

***Scytonostroma portentosum* (Berk. & M.A. Curtis) Donk**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003)

***Scytonostroma rhizomorpharum* S.S. Rattan**

Distribution: Roraima (Jesus 1996)

***Meruliaceae* Rea**

***Aquascypha hydrophora* (Berk.) D.A. Reid**

Distribution: Amapá [Batista et al. (1966), as *Stereum hydrophorum* Berk.]; Amazonas [Hennings (1904), as *Hymenochaete crateriformis* Henn.; Burt (1920), as *S. hydrophorum*; Reid (1965)]; Pará [Silva & Minter (1995), as *S. Hydrophorum*; Sotão et al. (2008)]

***Cymatoderma caperatum* (Berk. & Mont.) D.A. Reid**

Distribution: Amazonas [Hennings (1904), Theissen (1912), as *Thelephora caperata* Berk. & Mont.]; Pará (Theissen 1912); Roraima (Jesus 1996)

***Cymatoderma dendriticum* (Pers.) D.A. Reid**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003); Amazonas [Hennings (1904), as *Cladoderris dendritica* (Pers.) Berk.; Theissen (1912), as *Cladoderris crassa* (Klotzsch) Fr.; Reid (1965)]; Pará (Reid 1965); Rondônia [Capelari & Maziero (1988), as *C. dendritica*]; Roraima (Jesus 1996)

***Cotylidia aurantiaca* (Pers.) A.L. Welden**

Distribution: Amazonas (Reid 1965)

***Flaviporus liebmannii* (Fr.) Ginns**

Distribution: Amazonas [Hennings (1904), as *Polystictus stereinus* (Berk. & M.A. Curtis) Sacc.]; Pará [Sotão et al. 2008]; Rondônia [Capelari & Maziero (1988), as *Antrodiella liebmannii* (Fr.) Ryvarden]

***Gloeoporus dichrous* (Fr.) Bres.**

Distribution: Amazonas [Theissen (1911), as *Polyporus dichrous* Fr.]

***Gloeoporus cystidiatus* Ryvarden**

Distribution: Amazonas (Ryvarden 1987)

***Gloeoporus thelephoroides* (Hook.) Cunningham**

Distribution: Mato Grosso [Theissen (1911), Fidalgo (1968), as *Gloesporus conchooides* Mont.]; Pará [Hennings (1908), Theissen (1911), as *G. conchooides*; Sotão et al. (2008)]

***Hypochnicium lundellii* (Bourdot) J. Erikss.**

Distribution: Roraima (Jesus 1996)

***Hypochnicium punctulatum* (Cooke) J. Erikss.**

Distribution: Acre [Bononi (1992), as *Hypochnicium ounctulatum* (Cooke) Erikss.]

***Irpea lacteus* (Fr.) Fr.**

Distribution: Pará (Sotão et al. 2008)

***Junghuhnia undigera* (Berk. & M.A. Curtis) Ryvarden**

Distribution: Pará (Sotão et al. 2003)

***Rigidoporus amazonicus* Ryvarden**

Distribution: Amazonas (Ryvarden 1987)

***Rigidoporus biokoensis* (Bres. ex Lloyd) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Rigidoporus lineatus* (Pers.) Ryvarden**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003)

***Rigidoporus microporus* (Fr.) Overeem**

Distribution: Acre [Silva & Minter (1995), as *Fomes lignosus* Klotz]; Amazonas [Hennings (1940), as *Fomes auberianus* (Mont.) Murrill; Theissen (1911), as *Polyporus lignosus* Klotzsch; Torrend (1924, 1938)]; Pará [Batista et al. (1966), Silva & Minter (1995), as *F. lignosus*; Hennings (1908), as *F. auberianus*; Sotão et al. 2008]; Rondônia (Capelari & Maziero 1988); Roraima (Jesus 1996)

***Rigidoporus ulmarius* (Sowerby) Imazeki**

Distribution: Pará (Sotão et al. 2008)

***Rigidoporus vinctus* (Berk.) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Podoscypha brasiliensis* D.A. Reid**

Distribution: Acre (Bononi 1992); Pará (Reid 1965)

***Podoscypha bubalina* D.A. Reid**

Distribution: Amazonas (Reid 1965)

***Podoscypha elegans* (G. Mey.) Pat.**

Distribution: Amazonas [Hennings (1904), Theissen (1912), as *Stereum elegans* (G. Mey.) Fr.]

***Podoscypha fulvonitens* (Berk.) D.A. Reid**

Distribution: Amazonas (Reid 1965)

***Podoscypha ovalispora* D.A. Reid**

Distribution: Amazonas (Reid 1965)

***Podoscypha petalodes* subsp. *petalodes* (Berk.) Boidin**

Distribution: Amazonas [Hennings (1904), as *Stereum petalodes* Berk.]

***Podoscypha tomentipes* (Overh.) D.A. Reid**

Distribution: Amazonas (Reid 1965)

***Steccherinum ochraceum* (Pers.) Gray**

Distribution: Rondônia (Capelari & Maziero 1988)

***Steccherinum reniforme* (Berk. & M.A. Curtis) Banker**

Distribution: Amazonas (Bononi 1981); Mato Grosso (Hjortstam & Bononi et al. 1986)

***Stereopsis radicans* (Berk.) D.A. Reid**

Distribution: Acre (Bononi 1992); Mato Grosso [Theissen (1912), as *Thelephora radicans* Berk.]

***Stereopsis hiscens* (Berk. & Ravenel) D.A. Reid**

Distribution: Amazonas [Reid (1965), as *Stereopsis hiscens* var. *macrospora* D.A. Reid]

Meripilaceae Jülich

***Hydnopolyporus fimbriatus* (Fr.) D.A. Reid**

Distribution: Amazonas [Hennings (1904), as *Polyporus warmingii* Berk.; Theissen (1911), Batista et al. (1966), Silva & Minter (1995), as *Polystictus fimbriatus* Fr.]; Rondônia (Capelari & Maziero 1988)

Peniophoraceae Lotsy

***Peniophora albobadia* (Schwein.) Boidin**

Distribution: Pará [Sotão et al. (2003), Campos et al. (2005), as *Stereum albobadium* (Schwein.) Fr.]

Polyporaceae Corda

***Abundisporus roseoalbus* (Jungh.) Ryvarden**

Distribution: Pará [Sotão et al. (2008), as *Abundisporus subflexibilis* (Berk. & M.A. Curtis) Par.]

***Cinereomyces lenis* (P. Karst.) Spirin**

Distribution: Pará [Sotão et al. (2008), as *Skeletocutis lenis* (P. Karst.) Niemelä]

***Coriolopsis byrsina* (Mont.) Ryvarden**

Distribution: Acre (Bononi 1992)

***Coriolopsis floccosa* (Jungh.) Ryvarden**

Distribution: Pará [Sotão et al. (2002a, 2003), Martins-Júnior (2007), as *C. rigida* (Berk. & Mont.) Murrill; Sotão et al. (2002b), as *Trametes rigida* Berk. & Mont.; Sotão et al. (2008)]; Roraima [Jesus (1996), as *C. rigida*]

***Datronia caperata* (Berkeley) Ryvarden**

Distribution: Acre (Bononi 1992); Amapá [Batista et al. (1966), Silva & Minter (1995), as *Polystictus carperatus* Berk.]; Mato Grosso [Theissen (1911), as *P. carperatus*]; Pará [Batista et al. (1966), Silva & Minter (1995), Sotão et al. (2002b), as *Trametes caperata* (Berk.) Teixeira; Sotão et al. (2008)]; Rondônia (Capelari & Maziero 1988)

***Datronia mollis* (Sommerf.) Donk**

Distribution: Acre (Bononi 1992)

***Datronia stereoides* (Fr.) Ryvarden**

Distribution: Acre (Bononi 1992); Amapá [Batista et al. (1966), Silva & Minter (1995), as *Polystictus stereoides* Fr.]; Rondônia (Capelari & Maziero 1988); Pará [Silva & Minter (1995), as *P. stereoides*]

***Daedaleopsis confragosa* (Bolton) J. Schröt**

Distribution: Pará [Batista et al. (1966), as *Daedalea confragosa* Pers.]

***Diplomitoporus allantosporus* Ryvarden & Iturriaga**

Distribution: Pará (Martins-Júnior et al. 2008)

***Earliella scabrosa* (Pers.) Gilbn. & Ryvarden**

Distribution: Acre (Bononi 1992); Amapá [Sotão et al. (1991), as *E. corrugata* (Pers.) Murrill; Sotão et al. (2002a)]; Amazonas [Hennings (1904), as *Polystictus persoonii* (Mont.) Cooke]; Pará (Sotão et al. 1997, 2002b, 2008); Roraima (Jesus 1996); Rondônia (Capelari & Maziero 1988)

***Echinochaete brachypora* (Mont.) Ryvarden**

Distribution: Pará (Sotão et al. 1997, 2002b, 2008)

***Favolus tenuiculus* P. Beauv.**

Distribution: Acre (Bononi 1992), as *Favolus tessellatus* Mont.; Amazonas [Hennings (1904), as *F. paraguayensis* Speg. and *Favolus brasiliensis* Fr.; Theissen (1911), as *F. brasiliensis* Fr.]; Pará [Batista et al. (1966), Silva & Minter (1995), as *F. brasiliensis* and *F. paraguayensis*; Sotão et al. (1997, 2002b), as *Polyporus tenuiculus* (P. Beauv.) Fr.; Sotão et al. (2008)]; Rondônia [Capelari & Maziero (1988), as *F. brasiliensis*]

***Flabellophora obovata* (Jungh.) Núñez & Ryvarden**

Distribution: Amazonas [Hennings (1904), Theissen (1911), Silva & Minter (1995), Batista et al. (1966), as *Polystictus mutabilis* (Berk. & M.A. Curtis) Cooke]; Pará (Sotão et al. 1997, 2002b, 2008; Martins-Júnior 2007); Rondônia [Capelari & Maziero (1988), as *P. mutabilis*]

***Fomes fasciatus* (Schw.) Cooke**

Distribution: Pará (Sotão et al. 2008); Roraima (Jesus 1996)

***Funalia polyzona* (Pers.) Niemelä**

Distribution: Acre (Bononi 1992); Amapá [Batista et al. (1966), Silva & Minter (1995), as *Polystictus occidentalis* (Klotzsch) Fr.]; Amazonas [Henning (1904), as *P. occidentalis*]; Pará [Sotão et al. (2002b), as *Trametes polyzona* (Pers.) Corner; Sotão et al. (2008), as *Coriolopsis polyzona* (Pers.) Ryvarden]

***Grammothele fuligo* (Berk. & Br.) Ryvarden**

Distribution: Pará [Henning (1908), as *Polyporus buttneri* Henn.; Batista et al. (1966); Silva & Minter (1995)]; Roraima (Jesus 1996)

***Grammothele lineata* Berk. & M.A. Curtis**

Distribution: Roraima (Jesus 1996)

***Grammothele subargentea* (Speg.) Rajchenb.**

Distribution: Roraima (Jesus 1996)

***Hexagonia hydnoides* (Fr.) Fidalgo**

Distribution: Amazonas [Henning (1904), as *Polystictus hydnoides*; Theissen (1911), Batista et al. (1966), Silva & Minter (1995), as *Trametes hydnoides* (Sw.) Fr.]; Mato Grosso (Fidalgo 1968); Pará [Sotão et al. 2002a, 2002b, 2008; Campos et al. 2005]; Rondônia (Capelari & Maziero 1988)

***Hexagonia variegata* Berk.**

Distribution: Amazonas [Henning (1904); Theissen 1911]; Pará [Sotão et al. (2003), Campos et al. (2005), as *Hexagonia papyraceae* Berk.; Sotão et al. (2008)]; Roraima [Jesus (1996), as *H. papyraceae* Berk.]

***Lentinus copulatus* (Ehrenb.) Henn**

Distribution: Amazonas (Henning 1904)

***Lentinus crinitus* (L.) Fr.**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003); Amazonas [Henning (1904), as *L. subcervinus* Berk. & M.A. Curtis]; Mato Grosso (Fidalgo 1968); Pará (Sotão et al. 1997, 2002a, 2003, 2008; Campos et al. 2005); Rondônia (Capelari & Maziero 1988)

***Lentinus scleropus* (Pers.) Fr.**

Distribution: Amazonas [Henning (1904), as *Lentinus vellereus* Berk. & M.A. Curtis]

***Lentinus strigosus* (Schwein.) Fr.**

Distribution: Roraima [Jesus (1996), as *Panus rufus* Fr.]

***Lentinus velutinus* Fr.**

Distribution: Amazonas [Henning (1904), Batista et al. (1966), Silva & Minter (1995), as *Polystictus velutinus* Fr.]; Mato Grosso (Fidalgo 1968); Rondônia (Capelari & Maziero 1988); Pará [Silva & Minter (1995), as *P. velutinus*]

***Lentinus villosus* Klotzsch**

Distribution: Pará (Batista et al. 1966; Silva & Minter 1995)

***Lopharia cinerascens* (Schweinitz) G. Cunningham**

Distribution: Amazonas [Burt (1920), as *Stereum cinerascens* (Schwein.) Massee]; Mato Grosso [Theissen (1912), as *Lloydella cinerascens* (Schwein.) Bres.]; Pará [Hjortstam & Bononi (1986), Sotão et al. (2002a, 2003), Campos et al. (2005), as *S. cinerascens*; Sotão et al. (2008)]

***Loweporus tephroporus* (Mont.) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Megasporoporia cavernulosa* (Berk.) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Megasporoporia setulosa* (Henn.) Rajchenb.**

Distribution: Pará (Sotão et al. 2008)

***Microporellus dealbatus* Berk. & M.A. Curtis) Murrill**

Distribution: Amazonas (Silva & Minter 1995); Pará (Sotão et al. 2008); Rondônia (Capelari & Maziero 1988)

***Nigrofomes melanoporus* (Mont.) Murrill**

Distribution: Amazonas [Sousa (1980), as *Phellinus melanoporus* (Mont.) G. Cunn.]; Pará [Sousa (1980), as *P. melanoporus*; Sotão et al. (2008)]; Roraima [Sousa (1980), as *P. melanoporus*]; Rondônia [Sousa (1980), as *P. melanoporus*]

***Nigrohydnum nigrum* Ryvarden**

Distribution: Amazonas (Ryvarden 1987)

***Nigroporus rigidus* Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Nigroporus vinosus* (Berk.) Murrill**

Distribution: Acre (Bononi 1992); Amapá (Sotão et al. 1991, 2002a, 2003); Rondônia (Capelari & Maziero 1988)

***Pachykytospora alabamae* (Berk. & Cooke) Ryvarden**

Distribution: Pará (Sotão et al. 2008); Roraima (Jesus 1996)

***Perenniporia aurantiaca* (A. David & Rajchenb.) Decock & Ryvarden**

Distribution: Pará (Decock & Ryvarden 1999; Sotão et al. 2008)

***Perenniporia contraria* (Berk. & M.A. Curtis) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Perenniporia detritus* (Berk.) Ryvarden**

Distribution: Amazonas [Decock & Ryvarden (1999), as *Perenniporia dedrita* (Berk.) Ryvarden]

***Perenniporia inflexibilis* (Berk.) Ryvarden**

Distribution: Pará (Sotão et al. 2008); Roraima [Jesus (1996), as *P. glaucopora* (Lloyd) Ryvarden]

***Perenniporia martii* (Berk.) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Perenniporia medulla-panis* (Fr.) Donk**

Distribution: Pará [Decock & Ryvarden (1999), as *Perenniporia chromatica* (Berk. & Broome) Decock & Ryvarden; Sotão et al. 2008]

***Perenniporia ohiensis* (Berk.) Ryvarden**

Distribution: Mato Grosso [Theissen (1911, as *Trametes ohiensis* Berk.); Pará (Sotão et al. 2008)]

***Perenniporia stipata* Ryvarden**

Distribution: Pará (Sotão et al. 2008); Roraima (Ryvarden 1987)

***Perenniporia sinuosa* Ryvarden**

Distribution: Amazonas (Ryvarden 1987)

***Phaeolus schweinitzii* (Fr.) Pat.**

Distribution: Acre (Bononi 1992)

***Polyporus arcularius* (Batsch) Fr.**

Distribution: Pará (Sotão et al. 2008); Rondônia (Capelari & Maziero 1988)

***Polyporus dictyopus* Mont.**

Distribution: Amazonas [Henning (1904), as *P. vernicosus* Berk.; Torrend (1924)]; Pará [Sotão et al. (1997), as *P. infernalis* Berk.; Sotão et al. (2002b, 2008)]; Rondônia [Capelari & Maziero (1988), as *Melanopus infernalis* (Berk.) Pat.]; Roraima [Jesus (1996), as *P. infernalis* Berk.]

***Polyporus gramocephalus* Berk.**

Distribution: Amazonas (Henning 1904, Batista et al. 1966, Silva & Minter 1995)

***Polyporus guianensis* Mont.**

Distribution: Acre (Bononi 1992; Torrend 1920); Amazonas (Batista et al. 1966, Silva & Minter 1995); Pará (Sotão et al. 1997, 2002b, 2008)

***Polyporus ianthinus* Gibertoni & Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Polyporus leprieurii* Mont.**

Distribution: Amazonas (Henning 1904; Theissen 1911); Acre (Bononi 1992); Mato Grosso [Fidalgo (1968), as *Melanopus leprieurii* (Mont.) Pat.]; Pará (Sotão et al. 1997, 2002b, 2008); Rondônia [Capelari & Maziero (1988), as *M. leprieurii*]

***Polyporus rugulosus* Lév.**

Distribution: Amapá [Batista et al. (1966), Silva & Minter (1995), as *Polyporus plebeius* Berk.]

***Polyporus tricholoma* Mont.**

Distribution: Amazonas [Henning (1904), as *Polyporus flexipes* Fr.; Batista et al. (1966); Teixeira & Fidalgo (1983); Silva & Minter (1995)]; Acre (Bononi 1992); Pará (Sotão et al. 2002b, 2008); Rondônia (Capelari & Maziero 1988)

***Poronidulus conchifer* (Schwein.) Murrill**

Distribution: Amapá [Sotão et al. (1991, 2002a, 2003), as *Trametes conchifer* (Schwein.) Fr.]

***Pseudofavolus cucullatus* (Mont.) Pat**

Distribution: Rondônia (Capelari & Maziero 1988)

***Pseudofavolus miquelianus* (Mont.) Pat.**

Distribution: Rondônia (Capelari & Maziero 1988)

***Pycnoporus sanguineus* (Fr.) Murrill**

Distribution: Amapá [Sotão et al. 1991, 1996, 2002a, 2003]; Amazonas [Hennings (1904), as *Polystictus sanguineus* (L.) G. Mey.]; Pará [Batista et al. (1966), Silva & Minter (1995), as *P. sanguineus*; Sotão et al. (1997, 2002a, 2002b, 2003, 2008); Campos et al. (2005); Rondônia (Capelari & Maziero 1988)]

***Pyrofomes lateritius* (Cooke) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Stiptophyllum erubescens* (Berk.) Ryvarden**

Distribution: Acre (Bononi 1992); Amazonas [Hennings (1908), as *Lentinus schomburgkianus* Henn.]; Mato Grosso [Fidalgo (1968), as *Lenzites erubescens* (Berk.) Sacc.]

***Tinctoporellus epimiltinus* (Berk. & Broome) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Trametes cotonnea* (Pat. & Har.) Ryvarden**

Distribution: Acre (Bononi 1992); Pará (Sotão et al. 2008)

***Trametes cubensis* (Mont.) Sacc.**

Distribution: Pará (Sotão et al. 2002b, 2008)

***Trametes elegans* (Spreng.) Fr.**

Distribution: Pará [Sotão et al. (2002a, 2003); Sotão et al. (2008), as *Lenzites elegans*]; Rondônia (Capelari & Maziero 1988)

***Trametes incerta* (Curr.) Cooke**

Distribution: Amazonas [Torrend (1935), as *Hexagonia erubescens* Berk.]

***Trametes lactinea* (Berk.) Sacc.**

Distribution: Pará (Sotão et al. 2008)

***Trametes marianna* (Pers.) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Trametes maxima* (Mont.) A. David & Rajchenb.**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003); Pará (Sotão et al. 2008)

***Trametes membranacea* (Sw.) Kreisel**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003); Pará (Sotão et al. 1997, 2002b, 2008)

***Trametes modesta* (Kunze ex Fr.) Ryvarden**

Distribution: Acre [Batista et al. (1966), Silva & Minter (1995), as *Polystictus modestus* Fr.; Bononi (1992)]; Amazonas [Hennings (1904), as *Polyporus albocervinus* Berk.]; Pará (Sotão et al. 2008); Rondônia (Capelari & Maziero 1988); Roraima (Jesus 1996)

***Trametes nivosa* (Berk.) Murrill**

Distribution: Amapá [Sotão et al. (1991, 2002a, 2003), as *Tyromyces chioneus* (Fr.) P. Karst.]; Pará [Sotão et al. (2002a, 2003), Campos et al. (2005), as *T. chioneus*; Sotão et al. (2008)]

***Trametes pubescens* (Schumach.) Pilát**

Distribution: Acre (Bononi 1992)

***Trametes sclerodepsis* Berk.**

Distribution: Rondônia (Capelari & Maziero 1988)

***Trametes versicolor* (L.) Lloyd**

Distribution: Pará [Batista et al. (1966), Silva & Minter (1995), as *Polystictus versicolor* (L.) Fr.]

***Trametes villosa* (Sw.) Kreisel**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003); Pará [Batista et al. (1966), Silva & Minter (1995), as *Polystictus piniarius* Fr.]; Roraima (Jesus 1996)

***Trichaptum byssogenum* (Jungh.) Ryvarden**

Distribution: Pará (Sotão et al. 2002a, 2003, 2008; Campos et al. 2005); Rondônia (Capelari & Maziero 1988); Roraima (Jesus 1996)

***Trichaptum perrottetii* (Lév.) Ryvarden**

Distribution: Amapá [Sotão et al. (1991, 2002a, 2003), as *T. trichomallum* (Berk. & Mont.) Murrill]; Pará (Sotão et al. 2008)

***Trichaptum sector* (Ehrenb.) Kreisel**

Distribution: Amazonas [Hennings (1904), Theissen (1911), Batista et al. (1966), Silva & Minter (1995), as *Polystictus floridanus* Speg.]; Pará [Batista et al. (1966), Silva & Minter (1995), as *Polystictus sector* Fr.; Sotão et al. (2003), Campos et al. (2005), as *Trichaptum abientinum* (Dicks.) Ryvarden; Sotão et al. (2002b, 2008)]; Rondônia (Capelari & Maziero 1988)

***Tyromyces singeri* Ryvarden**

Distribution: Amazonas (Ryvarden 1987)

***Phanerochaetaceae* Jülich**

***Ceriporiopsis flavidula* (Murrill) Ryvarden**

Distribution: Pará (Martins-Júnior 2007)

***Climacodon chlamydocystis* Maas Geest.**

Distribution: Mato Grosso (Bononi 1981)

***Climacodon pulcherrimus* (Berk. & M.A. Curtis) M.I. Nikol.**

Distribution: Pará, Rondônia (Bononi 1981)

***Inflatostereum glabrum* (Lév.) D.A. Reid**

Distribution: Amazonas [Hennings (1904), as *Stereum huberianum* Henn.; Reid (1965); Burt (1920), as *Stereum fissum* Berk.]; Pará (Sotão et al. 2008)

***Phanerochaete sordida* (P. Karst.) J. Erikss. & Ryvarden**

Distribution: Roraima (Jesus 1996)

***Schizophyllaceae* Quélet**

***Schizophyllum commune* Fr.**

Distribution: Amapá (Sotão et al. 1991, 2002a, 2003); Pará [Hennings (1908), Batista et al. (1966), Silva & Minter (1995), as *S. alneum* J. Schröt.; Sotão et al. (1997, 2002a, 2003); Campos et al. (2005)]; Rondônia (Capelari & Maziero 1988)

***Schizophoraceae* Jülich**

***Schizopora flavipora* (Berk. & M.A. Curtis ex Cooke) Ryvarden**

Distribution: Pará (Sotão et al. 2008)

***Stereaceae* Pilát**

***Aleurodiscus aurantius* (Pers.) J. Schröt.**

Distribution: Roraima (Jesus 1996)

***Stereum hirsutum* (Willd.) Pers.**

Distribution: Pará (Sotão et al. 2002a, 2003; Campos et al. 2005)

***Stereum ostrea* (Blume & T. Nees) Fr.**

Distribution: Mato Grosso [Burt (1920), as *Stereum fasciatum* (Schwein.) Fr.]; Pará [Theissen (1912), as *Stereum lobatum* (Kunze ex Fr.) Fr.; Batista et al. (1966), Silva & Minter (1995), as *Stereum australe* Lloyd; Sotão et al. (1997)]; Roraima [Jesus (1996), as *Stereum fasciatum* (Schwein.) Fr.]

Taxa excluded due to unresolved nomenclatural problems:

***Amauroderma angustum* (Berk.) Torrend**

Distribution: Amazonas (Torrend 1920, 1922)

***Amauroderma juruense* (Henn.) Torrend**

Distribution: Acre (Torrend 1922)

***Aphelaria aurantiaca* (Henn.) Corner**

Distribution: Amazonas (Silva & Minter 1995); Pará (Corner 1950)

***Auricularia delicata* (Fr.) Henn.**

Distribution: Mato Grosso (Fidalgo 1968); Pará (Batista et al. 1966); Rondônia (Capelari & Maziero 1988)

***Auricularia tremellosa* (Fr.) P. Henn.**

Distribution: Pará (Hennings 1908)

***Cantharellus helvelloides* Henn.**

- Distribution: Amazonas (Henning 1904)
- Ceraceomerulius rubicundus* (Litsch.) J. Erikss. & Ryvarden
 Distribution: Roraima (Jesus 1996)
- Clavaria scabra* Berk.
 Distribution: Amazonas (Corner 1950)
- Corticium abnorme* Henn.
 Distribution: Amazonas (Henning 1904)
- Corticium caesalpiniae* Henn.
 Distribution: Pará (Henning 1908)
- Fomes bomfimensis* Henn.
 Distribution: Amazonas (Henning 1904)
- Fomes cremeotomentosus* Henn.
 Distribution: Amazonas (Henning 1904)
- Fomes haskarlii* (Lév.) Bres.
 Distribution: Mato Grosso (Theissen 1911)
- Fomes personii* Fr.
 Distribution: Amazonas (Theissen 1911)
- Fomes scalaris* Berk.
 Distribution: Pará (Batista 1996)
- Fomes subamboinensis* Henn.
 Distribution: Acre (Batista 1966; Silva & Minter 1995); Amazonas (Henning 1904)
- Formitopsis albomarginata* var. *albomarginata* (Zipp. ex Lév.) Imazeki
 Distribution: Amazonas (Corner 1989)
- Ganoderma reniformis* Morgan
 Distribution: Pará (Batista et al. 1966; Silva & Minter 1995)
- Hexagonia glabra* (P. Beauv.) Ryvarden
 Distribution: Pará (Sotão et al. 2008)
- Hexagonia tenuis* J.M. Hook
 Distribution: Acre (Silva & Minter 1995); Amapá (Batista et al. 1966; Silva & Minter 1995); Amazonas (Henning 1904)
- Hymenochaete elegantissima* (Speg.) Speg.
 Distribution: Mato Grosso (Theissen 1912)
- Hypolyssus montagnei* Berk.
 Distribution: Amazonas (Henning 1904)
- Lachnocladium furcellatum* (Fr.) Pat.
 Distribution: Amazonas (Henning 1904, Theissen 1912); Pará (Theissen 1912)
- Lachnocladium hamatum* Henn.
 Distribution: Amazonas (Henning 1904)
- Lachnocladium madeirensense* Henn.
 Distribution: Amazonas (Henning 1904)
- Lachnocladium manaosense* Henn.
 Distribution: Amazonas (Henning 1905, Corner 1950)
- Lachnocladium strictissimum* Henn.
 Distribution: Amazonas (Henning 1904)
- Laschia papulata* Mont.
 Distribution: Acre (Silva & Minter 1995)
- Lentinus cladopus* Lév.
 Distribution: Amapá (Sotão et al. 1991, 2002a, 2003)
- Lentinus concavus* (Berk.) Corner
 Distribution: Acre (Bononi 1992)
- Lentinus lecomtei* Fr.
 Distribution: Pará (Henning 1908)
- Lenzites repanda* (Mont.) Fr.

- Distribution: Amapá (Batista et al. 1966; Silva & Minter 1995); Mato Grosso (Theissen 1911)
- Peniophora caesalpiniae* Henn.
Distribution: Pará (Henning 1908)
- Phanerochaete hydnoides* (Cooke & Mass.) Donk
Distribution: Mato Grosso (Bodoni 1981)
- Phellinus zealandicus* (Cooke) Teng
Distribution: Amazonas, Mato Grosso (Sousa 1980)
- Podoscypha nitidula* var. *nitidula* (Berk.) Pat.
Distribution: Amazonas (Reid 1965)
- Polyporus cachoeirasensis* Henn.
Distribution: Amazonas (Henning 1904)
- Polyporus cayenninus* Mont.: Fr.
Distribution: Rondônia (Capelari & Maziero 1988)
- Polyporus leprieurii* var. *juruanus* Henn.
Distribution: Amazonas (Henning 1904)
- Polyporus marmellosensis* Henn.
Distribution: Amazonas (Henning 1904)
- Polyporus sacer* var. *juruensis* Henn.
Distribution: Amazonas (Henning 1904)
- Polyporus sulfuratus* (Fr.) Trotter
Distribution: Mato Grosso (Theissen 1911)
- Polystictus bivalvis* Pers.
Distribution: Amazonas (Henning 1904)
- Polystictus hunteri* Lloyd
Distribution: Amazonas (Silva & Minter 1995); Pará (Batista et al. 1966)
- Polystictus radiatoscruposus* Henn.
Distribution: Henning (1905)
- Polystictus tenuis* Link ex Cooke
Distribution: Amapá (Batista et al. 1996); Pará (Silva & Minter 1995)
- Polystictus subochraceus* Lloyd
Distribution: Amapá (Batista et al. 1996); Pará (Silva & Minter 1995)
- Polystictus trichomallus* (Durieu & Mont.) Fr.
Distribution: Amazonas (Henning 1904); Pará (Batista et al. 1966; Silva & Minter 1995)
- Polystictus velutinus* Fr.
Distribution: Pará (Batista et al. 1996)
- Schizophyllum brasiliense* W.B. Cooke
Distribution: Pará (Sotão et al. 2002a, 2003)
- Schizophyllum umbrinum* Berk.
Distribution: Pará (Sotão et al. 2003)
- Steccherinum subochraceum* Bononi & Hjortstam
Distribution: Amapá (Henning (1908), Sotão et al. 1991, 2002a, 2003)
- Stereum molle* Berk.
Distribution: Mato Grosso (Theissen 1012)
- Stereum versicolor* (Sw.) Fr.
Distribution: Acre (Bononi 1992)
- Xerotus echinosporus* Henn.
Distribution: Amazonas (Henning 1904)

Excluded taxa due to non neo- or pantropical distribution

- Amauroderma fuscoporia* Wakef.
Record: Pará (Sotão et al. 2002b)

***Amauroderma rude* (Berk.) Torrend**

Distribution: Amazonas (Furtado 1981)

***Antrodia sinuosa* (Fr.) P. Karst.**

Record: Amapá (Sotão et al. 1991, 2002a, 2003); Pará (Sotão et al. 2003)

***Antrodiella canadensis* (Overh.) Niemelä**

Record: Roraima [Jesus (1996), as *A. overholtsii* Ryvarden & Gilb.]

***Fibricium rude* (P. Karst.) Jülich**

Record: Roraima (Jesus 1996)

***Ganoderma applanatum* (Pers.) Pat.**

Record: Amapá (Sotão et al. (1991, 2002a, 2003); Pará (Sotão et al. 1997, 2002a, 2002b, 2003)

***Ganoderma lucidum* (Mont.) Pat.**

Record: Amapá (Sotão et al. 1991, 2002a, 2002b, 2003); Mato Grosso (Theissen 1911, Fidalgo 1968); Pará (Sotão et al. 1997); Roraima (Jesus 1996)

***Gloeocystidiellum convolvens* (P. Karst.) Donk**

Record: Roraima (Jesus 1996)

***Hypodontia gossypina* (Parmasto) Hjortstam**

Record: Roraima [Jesus (1996), as *Fibrodonitia gossypina* Par.]

***Ischnoderma resinosum* (Schrad.) P. Karst.**

Record: Rondônia (Capelari & Maziero 1988)

***Junghuhnia crustacea* (Jungh.) Ryvarden**

Record: Acre (Bononi 1992); Pará (Sotão et al. 2002a, 2003)

***Lentinus badius* (Berk.) Berk**

Record: Roraima [Jesus (1996), as *Panus badius* (Berk.) Sing.]

***Lopharia papyrina* (Mont.) Boidin**

Record: Mato Grosso [Fidalgo (1968), as *Stereum papyrinum* Mont.]

***Lignosus sacer* (Afzel. ex Fr.) Ryvarden**

Record: Amazonas [Hennings (1904), as *Polystictus sacer* (Afzel. ex Fr.) Fr.]

***Mucronella calva* (Alb. & Schwein.) Fr.**

Record: Acre (Bononi 1992); Rondônia (Capelari & Maziero 1988)

***Phellinus crocatus* (Fr.) Ryvarden**

Record: Amazonas [Hennings (1904), as *Polystictus crocatus* (Fr.) Fr.]

***Phylloporia ribis* (Schumach.) Ryvarden**

Record: Amazonas, Pará, Rondônia [Sousa (1980), as *Phellinus ribis* (Schumach.) Quél.]

***Polyporus melanopus* (Pers.) Fr.**

Record: Amazonas (Hennings 1904)

***Polyporus rhizophilus* Pat.) Sacc.**

Record: Acre (Bononi 1992)

***Pycnoporus cinnabarinus* (Jacq.) Fr.**

Record: Amazonas [Silva & Minter (1995), as *P. cinnabarinus* var. *lactescens*]; Pará [Batista et al. (1966), as *P. cinnabarinus* (Jacq.) Fr.; Silva & Minter (1995), as *P. cinnabarinus* and *P. cinnabarinus* var. *Sanguinea*]

***Thelephora lutescens* Schwein.**

Record: Acre (Bononi 1992)

***Thelephora paraguaiensis* Corner**

Record: Acre (Bononi 1992); Rondônia (Capelari & Maziero 1988)

Acknowledgments

The authors would like to thank Dr. Gregory Mueller and Dr. Leif Ryvarden for critically reviewing of the manuscript and CNPq for the master scholarship of ACGS.

Literature cited

- Alexopoulos CJ, Mims CW, Blackwell M. 1996. Introductory Mycology. 4th ed., John Wiley and Sons, Inc., Nova York , 868p.
- Batista AC, Falcão RGS, Peres GEP, Moura NR. 1966. Fungi Paraenses (Revisão da Coleção de Paul C. Hennings, do Museu Paraense Emílio Goeldi). Instituto de Micologia. 506: 10–290.
- Bononi VL. 1981. Alguns Basidiomicetos hidnóides da região amazônica. *Rickia*. 9: 17–30.
- Bononi VL. 1992. Fungos macroscópicos de Rio Branco, Acre, Brasil. *Hoehnea*, 19(1/2): 31–37.
- Burt EA. 1920. The *Thelephoraceae* of North America XII. *Annals of the Missouri Botanical Garden*. 7: 81–238.
- Capelari M, Maziero R. 1988. Fungos macroscópicos do estado de Rondônia, região dos rios Jaru e Ji-Paraná. *Hoehnea*, 15: 28–36.
- Campos EL, Cavalcanti MAQ. 2000. Primeira ocorrência de *Phellinus mangroicus* (Imaz.) Imaz. para o Brasil. *Acta Botânica Brasileira*. 14(3): 263–265.
- Campos EL, Sotão HMP, Cavalcanti MAQ, Luz AB. 2005. Basidiomycetes de Manguezais da APA de Algodoal – Maiandrea, Pará, Brasil. *Boletim do Museu Paraense Emílio Goeldi*, sér. bot. 1: 141–146.
- Corner EJH. 1950. A monograph of Clavaria and allied genera. *Annals of Botany Memoirs*, 1: 740.
- Corner EJH. 1983. Ad. Polyporaceae I. Beih. *Zur. Nova Hedwigia*. 75: 5–182.
- Corner EJH. 1989. Ad Polyporaceas V. The genera *Albatrellus*, *Boletopsis*, *Coriolopsis* (dimitic), *Cristelloporia*, *Diacanthodes*, *Elmerina*, *Fomitopsis* (dimitic), *Gloeoporus*, *Grifola*, *Hapalopilus*, *Heterobasidion*, *Hydnopolyporus*, *Ischnoderma*, *Loweporus*, *Parmastomyces*, *Perenniporia*, *Pyrofomes*, *Stecchericum*, *Trechispora*, *Truncospora* and *Tyromyces*. *Beih Nova Hedwig* 96:1–218
- Decock C, Ryvarden L. 1999. Studies in neotropical polypores 1. Some coloured resupinate *Perenniporia* species. *Mycological Research* 103 (9): 1138–1144.
- Decock C, Ryvarden L. 1999. Studies in *Perenniporia dedrita* and its taxonomic synonyms. *Mycologia* 91(2): 386–395.
- Fidalgo MEPK. 1968. Contribution to the fungi of Mato Grosso, Brazil. *Rickia*, 3: 171–219.
- Furtado JS. 1967. Some tropical species of *Ganoderma* (Polyporaceae) with pale context. *Persoonia* 4: 379–389.
- Furtado JS. 1981. Taxonomy of *Amauroderma* (*Basidiomycetes*, *Polyporaceae*). *Memoirs of the New York Botanical Garden*, 34: 1–109.
- Gibertoni TB, Parmasto E, Cavalcanti MAQ. 2003. Non-poroid Hymenochaetaceae (Basidiomycota) of the Atlantic Rain Forest in Northeast Brazil - with a preliminary check list of Brazilian species. *Mycotaxon*. 87: 437–443.
- Gibertoni TB, Bernicchia A, Ryvarden L, Gomes-Silva AC. 2008. Bresadola s polypore collection at the Natural History Museum of Trento, Italy. 2. *Mycotaxon*. 104: 321–323.
- Gomes-Silva AC, Gibertoni TB, Ryvarden L. 2008. *Coltricia fragilissima* a new record to Brazil. *Mycotaxon*, 105: 469–472.
- Hennings P. 1904. Fungi Amazonici I. a cl. Ernesto Ule collecti. *Hedwigia*, 43(3): 154–186.
- Hennings P. 1905. Fungi Amazonici IV. a cl. Ernesto Ule collecti. *Hedwigia*, 44(2): 57–71.
- Hennings P. 1908. Fungi Paraensis III. *Hedwigia*. 48(1-2): 101–117.
- Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, Eriksson OE, Huhndorf S, James T, Kirk PM, Lücking R, Thorsten-Lumbsch H, Lutzoni F, Matheny PB, McLaughlin DJ, Powell MJ, Redhead S, Schoch CL, Spatafora JW, Stalpers JA, Vilgalys R, Aime MC, Aptroot A, Bauer R, Begerow D, Benny GL, Castlebury LA, Crous PW, Dai YC, Gams W, Geiser DM, Griffith GW, Gueidan C, Hawksworth DL, Hestmark G, Hosaka K, Humber RA, Hyde KD, Ironside JE, Köljalg U, Kurtzman CP, Larsson KH, Lichtwardt R, Longcore J, Miadlikowska J, Miller A, Moncalvo JM, Mozley-Standridge S, Oberwinkler

- F, Parmasto E, Reeb V, Rogers JD, Roux C, Ryvarden L, Sampaio JP, Schüßler A, Sugiyama J, Thorn RG, Tibell L, Untereiner WA, Walker C, Wang Z, Weir A, Weiss M, White MM, Winka K, Yao YJ, Zhang N. 2007. A higher-level phylogenetic classification of the Fungi. *Mycological Research* 111(5): 509–547.
- Hjortstam K, Bononi VL. 1986. Fungos corticióides do Brasil (Aphyllophorales). *Rickia*, 13: 113–125.
- Jesus MA. 1996. Contribution to the knowledge of wood-rotting fungi in Brazil. II. Checklist of fungi from Maraca Island, Roraima State. *Mycotaxon*, 57: 323–328.
- Léger JC. 1998. Le genre *Hymenochaete* léveillé. *Bibliotheca Mycologia*. 171: 1–139.
- Martins-Júnior AS. 2007. Basidiomycetes Poliporóides Xilófilos da Estação Científica Ferreira Penna, Pará, Brasil. 134p. Dissertação (Mestrado em Botânica) – Universidade Federal Rural da Amazônia - Museu Paraense Emílio Goeldi, Belém, Pará.
- Martins-Júnior AS, Gibertoni TB, Sotão HMP. 2008. *Diplomitoporus allantoporus* (*Basidiomycetes*): a new record to Brazil. *Mycotaxon*, 106: 195–198.
- Museu Paraense Emílio Goeldi. 2007. Amazônia: Fundamentos da ecologia da maior região de florestas tropicais. http://www.museu-goeldi.br/biodiversidade/o_amazonia.asp. Acesso em 06/06/2007.
- Reid DA. 1965. A monograph of the stipitate steroid fungi. *Beihefte zur Nova Hedwigia* 18: 1–184.
- Ryvarden, L. 1987. New and noteworthy polypores from tropical America. *Mycotaxon* 28: 525–541.
- Silva M, Minter DW. 1995. Fungi from Brasil - Recorded by Batista and co-workers. *Mycological Papers*, 169: 1–585.
- Sotão HMP, Bononi VLR, Figueiredo TS. 1991. *Basidiomycetes* de manguezais da Ilha de Maracá, Amapá, Brasil. *Boletim do Museu Paraense Emílio Goeldi*, sér. Bot., 7(1): 109–114.
- Sotão HMP, Figueiredo TS. 1996. Utilização do fungo *Pycnoporus sanguineus* (L.: Fr.) Murr. na cerâmica do Muruanum, Amapá. *Boletim do Museu Paraense Emílio Goeldi*, Série Botânica. 12(1): 15–20.
- Sotão HMP, Hennen JF, Gugliotta AM, Melo OA, Campos EL. 1997. Os fungos – *Basidiomycota*. In: Lisboa, P.L.B. (org.). Caxiuanã. Belém, Museu Paraense Emílio Goeldi. 213–219.
- Sotão HMP, Campos EL, Costa SPSEC, Melo AO, Azevedo JC. 2002a. Basidiomycetes macroscópicos de manguezais de Bragança, Pará, Brasil. *Hoehnea*. 29(3): 215–224.
- Sotão HMP, Gugliotta AM, Oliveira AP, Luz AB, Melo OA. 2002b. Fungos poliporóides. In: Lisboa, P.L.B. (org.). Caxiuanã, populações tradicionais, meio físico e diversidade biológica. Belém: Museu Paraense Emílio Goeldi. p. 433–444.
- Sotão HMP, CAMPOS EL, Gugliotta AM, Costa SPSEC. 2003. Fungos macroscópicos: *Basidiomycetes*. In: Fernandes, M. E. B. (org.). Os manguezais da costa norte brasileira. Maranhão: Fundação Rio Bacanga. 142 p.
- Sotão HMP, Gibertoni TB, Maziero R, Baseia I, Medeiros PS, Martins-Júnior A, Capelari M. 2008. Fungos macroscópios da Floresta nacional de Caxiuanã, Pará, Brasil: *Basidiomycota* (*Agaricomycetes*). In: Lisboa P.L.B. (org.). Caxiuanã: Desafios para conservação de uma Floresta Nacional na Amazônia. Belém: Museu Paraense Emílio Goeldi.
- Sousa MA. 1980. O gênero *Phellinus* Quélet (*Hymenochaetaceae*) na Amazônia brasileira. Tese de Doutorado, Instituto de Pesquisa da Amazônia/ Fundação Universidade do Amazonas, Manaus, 190p.
- Steyaert RL. 1962. Genus *Ganoderma* (Polyporaceae) taxa nova-2. *Bulletin du Jardin Botanique de L'etat* Bruxelles. 32(1): 89–104.
- Steyaert RL. 1980. Study of some *Ganoderma* species. *Bulletin du Jardin Botanique de L'etat* Bruxelles. 50: 135–186.

- Teixeira AR, Fidalgo O. 1983. *Polyporus tricholoma* Mont. (Polyporaceae). Revista Brasileira de Botânica. 6 (2): 129–132.
- Theissen F. 1911. *Polyporaceae austro-brasilienses imprimis rio grandenses*. Kaiserlich-Königlichen Hof- und Staatsdruckerei, Wien, 60p.
- Theissen F. 1912. *Hymenomycetes Riograndenses*. Brotéria, ser. bot. 10: 5–28.
- Torrend C. 1920. Les polyporacées du Brésil. *Polyporaceus stipitées*. Brotéria, Série Botânica. 18 (1): 21–42.
- Torrend C. 1922. Les polyporacées du Brésil. Les genres *Amauroderma* et *Lignosus*. Brotéria, Série Botânica. 18 (2): 21–42.
- Torrend C. 1924. Les polyporacées s du Brésil. II. Brotéria, Ser. Bot. 21(1): 12–42.
- Torrend C. 1926. Les polyporacées stipitées du Brésil IV. Brotéria, ser. bot. 22: 5–19.
- Torrend C. 1935. Les polyporacées du Brésil. Brotéria, Ser. Bot. 31(3): 108–120.
- Torrend C. 1938. As poliporáceas da Bahia e estados limítrofes. Anais da Primeira Reunião Sul-Americana de Botânica, vol. II: 326–341.