

## Contribution to the lignocellulolytic fungi (Basidiomycetes) of the Atlantic Rain Forest in Southern Brazil

CLAUDIA GROPOSO  
*calitagr@hotmail.com*

and

CLARICE LOGUERCIO-LEITE

*Laboratório de Micologia, Departamento de Botânica  
Universidade Federal de Santa Catarina, Florianópolis,  
Santa Catarina, Brazil*

**Abstract** – Forty-seven species of wood inhabiting Basidiomycetes are reported from the Atlantic Rain Forest of the Parque Estadual da Serra do Tabuleiro, Santa Catarina, Brazil. Fourteen are new records for the state. All the species are new records for the studied area.

**Key words:** wood rot Basidiomycetes, checklist, xilophilous.

### Introduction

The macrofungi have perhaps the longest history of diversity studies of any mycota. Nevertheless, they are understudied over most of the world. Taxonomic obstacles and the absence of long-term studies prevent conclusive answers even to basic questions about the number of species at a specific location or whether diversity is greater in one type of forest than in another (Lodge et al. 2004). It is known that the species of lignocellulolytic basidiomycetes are extremely abundant in all forest types and that they are the major wood decomposers in most ecosystems (Fryar et al. 1999). These fungi have received special attention of researchers in the last decades due to their potential applications in pollutant purification, soil bioremediation, and antibiotic production (Blanchette 1995; Kotterman et al. 1994; Smânia et al. 2003). However, in many geographical areas, especially in the tropical forest systems, it is clear that the known number of described fungal species represents only a small fraction of the number believed to exist there (Gilbert & Sousa 2002). The plant flora is better known than the fungi both in the Neotropical area (Thomas 1999) and the Brazilian Atlantic Forest. The latter is particularly relevant for studies because of its high biological diversity and severely threatened status (Tabanez & Viana 2000). An increased understanding of fungal diversity and natural history of fungi will contribute to the knowledge of the local biota and will greatly strengthen initiatives to protect and use sustainably our natural resources (Rossman et al. 1998).

According to available records, fungi were first collected in Santa Catarina State in 1815 by Adalberto de Chamisso, and only sporadically thereafter until Möller's activity in 1890 (Loguercio-Leite 1991). Starting in 1986, surveys were periodically conducted by researchers of the Mycology Laboratory (BOT/CCB/UFSC), who specifically targeted polyporoid lignocellulolytic fungi. Since then several studies have been published with data on collections from the Atlantic Forest, Santa Catarina Island, and other areas in the state. This has produced approximately 60 new species

records for the state, six of which are new to science (Loguercio-Leite & Wright 1991, 1998; Loguercio-Leite et al. 1998, 2001, 2002).

### **Material and Methods**

The Atlantic Forest is composed of two major vegetation types: the Atlantic Rain Forest and the Atlantic Semi-deciduous Forest (Morellato & Haddad 2000). The state of Santa Catarina (26°-30° S lat, 48°30'-54° W long) in Southern Brazil currently has only 17.4% of its original area covered by the Atlantic Forest (Ministério do Meio Ambiente 2002).

The present study was conducted in Parque Estadual da Serra do Tabuleiro, which belongs to the Atlantic Rain Forest region. The Parque (27° 42' 09"-28° 34' 09" S lat, 48° 57' 23"-48° 43' 09" W long), which covers 900 km<sup>2</sup> and is located at the center-east portion of the state of Santa Catarina, includes nine municipalities (Fig. 1): Palhoça, Santo Amaro da Imperatriz, Águas Mornas, São Bonifácio, Imaruí, Garopaba, Paulo Lopes, São Martinho and Florianópolis (Batista 2003).

Highly diverse, the Parque Estadual da Serra do Tabuleiro contains representatives of the different landscapes types of Santa Catarina. The diverse ecosystems and habitats support known rare, endangered and endemic species, with many new species yet to be recorded. Therefore, this area both serves as a highly important genetic reservoir and offers opportunities for biodiversity and conservation research as well (Rosário 2003).

Periodical fungal inventories were conducted between July 2000 and July 2001. Basidiomes were studied using macroscopic (e.g., size, colour, number pores/mm, length of tubes) and microscopic (presence/absence of structures, dimensions) vegetative and reproductive characters (Ryvarden 1991, Singer 1975). Measurements and drawings were made from slide preparations stained with 1% aqueous phloxine and 5% KOH. Specimens were identified to species using specialized references and comparing collections with FLOR, BAFC, ICN and PACA Herbaria collections. Vouchers were deposited in FLOR Herbarium (Universidade Federal de Santa Catarina) after examination. (Holmgren et al. 1990). Nomenclature and authorities are from Kirk et al. (2001).

### **Results and Discussion**

In the state of Santa Catarina, 163 species of lignocellulolytic fungi (Basidiomycetes) have been recorded since 1986. In the investigated area 47 species were found, representing about 28 percent of the total number of polypores reported for the State, until now. Those species are distributed in 29 genera and 11 families. Fourteen of the identified species are new records for the State and are marked with \* in the checklist. All species are new citations for the area.

As this study was based on collecting during a period of 13 months and, as some fungi form its basidiomes irregularly, many species were probably unrecorded. Mycelia belonging to a species with low productivity may exist in the substrate over many years but only rarely produce basidiomes except under optimal conditions (Straatsma & Krisai-Greilhuber 2003)

Thirty-two (68%) of the species collected in the Parque Estadual da Serra do Tabuleiro, show a tropical (neotropical and pantropical) distribution, while four

[*Amauroderma corneri* Gulaid & Ryvarden, *Antrodiella multipileata* L. Leite & Wright, *Hemingsia brasiliensis* (Speg.) Speg., *Skeletocutis roseolus* (Rick ex Theissen) Rajch.] are known only from Brazil. Fifteen (32%) species are cosmopolitan.

The genera with the largest number of species (4) were *Ganoderma* and *Phellinus*. *P. grenadensis* was most frequent (8), mostly on trunks of *Piptadenia gonoacantha*. However, host data are poor in the tropics, particularly as the great diversity of species makes host identification difficult, particularly when basidiomes fruit on dead trees or fallen trunks (Gilbert et al. 2002).

This study substantially increases the geographic ranges of the lignocellulolytic fungi. Obviously more work is required to increase our knowledge on these fungi at the Park, an area of extreme biological importance.

#### - Checklist for the species at Parque Estadual da Serra do Tabuleiro:

##### AGARICALES

###### Tricholomataceae

\**Lentinus crinitus* (L.) Fr.

Syst. Orb. Veg. p. 77. 1825.

BASIONYM: *Agaricus crinitus* L., 1763.

DISTRIBUTION: neotropical.

Ref.: Dennis, 1970; Guzmán & Johnson, 1974; Pavlich, 1976; Jejelowo & Abraham, 1998.

\**Lentinus strigosus* (Schw.) Fr.

Syst. Orb. Veg. p. 77. 1825.

BASIONYM: *Agaricus strigosus* Schw., 1822.

DISTRIBUTION: pantropical.

Ref.: Rick, 1938; Dennis, 1970; Hongo, 1974; Guzmán & Johnson, 1974; Pavlich, 1976; Pegler, 1987; Saber, 1997.

\**Panellus pusillus* (Pers. ex Lév.) Burdsall & Miller.

Nova Hedwigia Beih. 51: 85, 1975.

BASIONYM: *Gloeoporus pusillus* Pers. ex Lév., 1844.

DISTRIBUTION: cosmopolitan.

Ref.: Burdsall & Miller, 1975.

###### Schizophyllaceae

\**Schizophyllum commune* Fr.

Syst. Mycol. 1: 330, 1821.

DISTRIBUTION: cosmopolitan.

Ref.: Cooke, 1961.

##### HYMENOGYALES

###### Hymenochaetaceae

*Cyclomyces iodinus* (Mont.) Pat.

Essai Tax. p. 98, 1900.

BASIONYM: *Polyporus iodinus* Mont., 1841.

DISTRIBUTION: neotropical.

Ref.: Gerber, 1996.

\**Cyclomyces tabacinus* (Mont.) Pat.

Essai Tax. p. 98, 1900.

BASIONYM: *Polyporus tabacinus* Mont., 1835.

DISTRIBUTION: pantropical.

- Ref.: Ryvarde & Johansen, 1980; Nuñez & Ryvarde, 2000a.
- \*Hymenochaete corrugata* (Fr.) Lév.  
Ann. Sci. Nat. Bot. III 5: 152, 1846.  
BASIONYM: *Thelephora corrugata* Fr., 1815.  
DISTRIBUTION: cosmopolitan.  
Ref.: Job, 1985a,b; Dueñas & Tellería, 1988; Soares & Gugliotta, 1998; Dai, 2000.
- \*Hymenochaete minuscula* Cunn.  
Trans. Royal Soc. New Zeal. 85: 48, 1957.  
DISTRIBUTION: pantropical.  
Ref.: Job, 1985a; Azevedo & Guerrero, 1993.
- \*Hymenochaete sallei* Berk. & Curt.  
Linn. Soc. Bot. J. 10: 333, 1868.  
DISTRIBUTION: cosmopolitan.  
Ref.: Dennis, 1970; Job, 1985a; Azevedo & Guerrero, 1993; Zhang, 1999.
- Phellinus gilvus* (Schw.: Fr.) Pat.  
Ess. Tax. Hym. p. 97, 1900.  
BASIONYM: *Boletus gilvus* Schw., 1822.  
DISTRIBUTION: cosmopolitan.  
Ref.: Loguercio-Leite & Wright, 1995.
- Phellinus grenadensis* (Murr.) Ryv.  
Norw. J. Bot. 18: 234, 1972.  
BASIONYM: *Pyropolyporus grenadensis* Murr., 1908.  
DISTRIBUTION: pantropical.  
Ref.: Larsen & Cobb-Pouille, 1990; Gerber & Loguercio-Leite, 2000.
- Phellinus umbrinellus* (Bres.) Herrera & Bond.  
Mikol. Fitopatol. 14(1): 8, 1980.  
BASIONYM: *Poria umbrinella* Bres., 1896.  
DISTRIBUTION: pantropical.  
Ref.: Gilbertson & Ryvarde, 1987; Loguercio-Leite & Wright, 1995.
- Phellinus wahlbergii* (Fr.) Reid.  
Contr. Bolus. Herb. 7: 97, 1975.  
BASIONYM: *Trametes wahlbergii* Fr., 1848.  
DISTRIBUTION: pantropical.  
Ref.: Ryvarde & Johansen, 1980; Gilbertson & Ryvarde, 1987; Loguercio-Leite & Wright, 1995.
- Phylloporia spathulata* (Hook.) Ryv.  
Synop. Fung. 5: 196, 1991.  
BASIONYM: *Polyporus spathulatus* Hook., 1822.  
DISTRIBUTION: pantropical.  
Ref.: Ryvarde & Johansen, 1980.; Gerber, 1996; Wagner & Ryvarde, 2002.

## POLYPORALES

### Corticaceae

- Gloeoporus dichrous* (Fr.) Bres.  
Annls. Mycol. 14: 230, 1916.  
BASIONYM: *Polyporus dichrous* Fr., 1815.  
DISTRIBUTION: cosmopolitan.  
Ref.: Gilbertson & Ryvarde, 1986; Nuñez & Ryvarde, 2000b.

### Ganodermataceae

- Amauroderma camerarium* (Berk.) Furtado  
Revisão do gênero *Amauroderma* (tese): 140, 1968.  
BASIONYM: *Polyporus camerarius* Berk., 1856.

DISTRIBUTION: neotropical.

Ref.: Furtado, 1981.

*Amauroderma corneri* Gulaid & Ryvarden

Mycologia Helvetica 10(1): 28, 1998.

DISTRIBUTION: neotropical (known only Brazil).-

Ref.: Gulaid & Ryvarden, 1998.

*Ganoderma applanatum* (Pers.) Pat.

Bull. Soc. Mycol. France 5: 67, 1889.

BASIONYM: *Boletus applanatus* Pers., 1799.

DISTRIBUTION: cosmopolitan.

Ref.: Steyaert, 1975; Silveira & Guerrero, 1991; Leonard, 1998; Nuñez & Ryvarden, 2000a.

*Ganoderma resinaceum* Boudier.

Bull. Soc. Mycol. France 5: 72, 1889.

BASIONYM: *Ganoderma chaffangeonii* Pat., 1889.

DISTRIBUTION: cosmopolitan.

Ref.: Steyaert, 1972; Nuñez & Ryvarden, 2000a.

*Ganoderma tornatum* (Pers.) Bres.

Hedwigia 53: 55, 1912.

BASIONYM: *Polyporus tornatus* Pers., 1827.

DISTRIBUTION: pantropical.

Ref.: Leonard, 1998; Nuñez & Ryvarden, 2000a.

### Hapalopilaceae

*Bjerkandera adusta* (Willd.: Fr.) Karst.

Medd. Soc. Fauna Flora Fenn. 5: 38, 1879.

BASIONYM: *Boletus adustus* Willd., 1787.

DISTRIBUTION: cosmopolitan.

Ref.: Gilbertson & Ryvarden, 1986; Nuñez & Ryvarden, 2000b.

*Ceriporia mellea* (Berk. & Br.) Ryv.

Bull. Jard. Bot. Nat. Belg. 48: 98, 1978.

BASIONYM: *Poria mellea* (Berk. & Br.) Sacc., 1888.

DISTRIBUTION: pantropical.

Ref.: Nuñez & Ryvarden, 2000b.

### Meripilaceae

*Henningsia brasiliensis* (Speg.) Speg.

Bol. Acad. Nac. Ci. Córdoba 23: 411, 1919.

BASIONYM: *Polyporus brasiliensis* Speg., 1889.

DISTRIBUTION: neotropical (known only Brazil).

Ref.: Ginns, 1979; Gerber & Loguercio-Leite, 2000.

*Rigidoporus microporus* (Fr.) Overeem.

Icon. Fung. Malayensum 5: 1, 1924.

BASIONYM: *Polyporus microporus* Fr., 1821.

DISTRIBUTION: pantropical.

Ref.: Gilbertson & Ryvarden, 1986; Gugliotta, 1997.

*Rigidoporus ulmarius* (Sow.: Fr.) Imazeki.

Bull. Govt Forest. Exp. Stn. Meguro 57: 97, 1952.

BASIONYM: *Boletus ulmarius* Sow., 1797.

DISTRIBUTION: cosmopolitan.

Ref.: Silveira & Guerrero, 1991.

*Rigidoporus vinctus* (Berk.) Ryv.

Norw. J. Bot. 19: 139-144, 1972.

BASIONYM: *Polyporus vinctus*, Berk., 1852.

DISTRIBUTION: pantropical.

Ref.: Ryvarde, 1972; Setliff, 1972.

### **Polyporaceae**

*Earliella scabrosa* (Pers.) Gilbn. & Ryv.

Mycotaxon 22: 364, 1985.

BASIONYM: *Polyporus scabrosus* Pers., 1827.

DISTRIBUTION: neotropical.

Ref.: Gilbertson & Ryvarde, 1986; Gerber & Loguercio-Leite, 2000; Nuñez & Ryvarde, 2000b.

*Hexagonia papyracea* Berk.

Ann. Mag. Nat. Hist. 10: 379, 1843.

DISTRIBUTION: pantropical.

Ref.: Loguercio-Leite, 1994.

*Microporellus obovatus* (Jungh.) Ryv.

Norw. J. Bot. 19: 232, 1972.

BASIONYM: *Polyporus obovatus* Jungh., 1838.

DISTRIBUTION: pantropical.

Ref.: Mitra, 1999.

*Perenniporia subannosa* (Bres.) C. Decock, Herrera & Ryv.

Mycologia 93(1): 196-204, 2001.

BASIONYM: *Fomes subannosus* Bres., 1926.

DISTRIBUTION: neotropical.

Ref.: Gerber et al., 1999; Decock et al., 2001.

\**Polyporus arcularius* (Batsch) Fr.

Syst. Mycol. 1: 342, 1821.

BASIONYM: *Boletus arcularius* Batsch, 1783.

DISTRIBUTION: pantropical.

Ref.: Ryvarde & Johansen, 1980; Gilbertson & Ryvarde, 1986; Nuñez & Ryvarde, 2000b.

*Polyporus dictyopus* Mont.

Annls. Sci. Nat. Bot. II 3: 345, 1835.

DISTRIBUTION: pantropical.

Ref.: Ryvarde & Johansen, 1980; Loguercio-Leite, 1992.

*Polyporus tenuiculus* (Beauv.) Fr.

Syst. Mycol. 1: 344, 1821.

BASIONYM: *Favolus tenuiculus* Beauv., 1806.

DISTRIBUTION: pantropical.

Ref.: Loguercio-Leite, 1992.

*Pycnoporus sanguineus* (L.: Fr.) Murr.

Bull. Torrey Bot. Club 31: 421, 1904.

BASIONYM: *Boletus sanguineus* L., 1763.

DISTRIBUTION: pantropical.

Ref.: Nuñez & Ryvarde, 2000b.

\**Skeletocutis roseolus* (Rick ex Theissen) Rajch.

Nord. J. Bot. 7(5): 561, 1987.

BASIONYM: *Polystictus roseolus* Rick ex Theissen, 1911.

DISTRIBUTION: neotropical (known only Brazil).

Ref.: David, 1982; Rajchenberg, 1987; Silveira & Guerrero, 1991.

\**Trametes membranacea* (Swartz) Kreisel.

Monogr. Cienc. Univ. Habana 4 16: 83, 1971.

BASIONYM: *Boletus membranaceus* Swartz, 1806.

DISTRIBUTION: neotropical.

Ref.: Silveira & Guerrero, 1991.

*Trametes versicolor* (L.) Pilát.

Atlas Champ. Eur. Polypor. B 3: 261, 1939.

BASIONYM: *Boletus versicolor* L., 1753.

DISTRIBUTION: cosmopolitan.

Ref.: Silveira & Guerrero, 1991; Loguercio-Leite, 1993.

*Trametes villosa* (Swartz) Kreisel.

Monogr. Cienc. Univ. Habana 4 16: 83, 1971.

BASIONYM: *Boletus villosus* Swartz, 1806.

DISTRIBUTION: neotropical.

Ref.: Loguercio-Leite, 1993.

*Trichaptum sector* (Ehrenb.) Kreisel.

Rev. Gen. Pl. 3: 497, 1971.

BASIONYM: *Boletus sector* Ehrenb., 1820.

DISTRIBUTION: neotropical.

Ref.: Silveira & Guerrero, 1991; Gerber, 1996.

*Tyromyces leucomallus* (Berk. & Curt.) Murr.

North American Flora 9: 36, 1907.

BASIONYM: *Polyporus leucomallus* Berk. & Curt., 1868.

DISTRIBUTION: cosmopolitan.

Ref.: Gilbertson & Ryvarden, 1986; Gonçalves & Loguercio-Leite, 2001.

### **Steccherinaceae**

*Antrodiella multipileata* L. Leite & Wright.

Mycotaxon 41: 167, 1991.

DISTRIBUTION: neotropical.

Ref.: Loguercio-Leite & Wright, 1991.

*Antrodiella semisupina* (Berk. & Curt.) Ryv.

Prel. Polyp. Fl. East Africa p. 261, 1980.

BASIONYM: *Polyporus semisupinus* Berk. & Curt., 1872.

DISTRIBUTION: cosmopolitan.

Ref.: Gilbertson & Ryvarden, 1986; Nuñez & Ryvarden, 2000b.

*Flaviporus brownei* (Humb.) Donk.

Persoonia 1: 189, 1959.

BASIONYM: *Boletus brownii* Humb., 1793.

DISTRIBUTION: cosmopolitan.

Ref.: Ginns, 1980; Gerber & Loguercio-Leite, 1997; Gonçalves & Loguercio-Leite, 2001.

## **RUSSULALES**

### **Hericiaceae**

\**Laxitextum bicolor* (Pers.) Lentz.

Agric. Monogr. 24: 19, 1955.

BASIONYM: *Thelephora bicolor* Pers., 1801.

DISTRIBUTION: cosmopolitan.

Ref.: Eriksson & Ryvarden, 1976.

### **Stereaceae**

*Cymatoderma dendriticum* (Pers.) Reid.

Kew Bull. 1958: 523, 1959.

BASIONYM: *Thelephora dendritica* Pers., 1827.

DISTRIBUTION: pantropical.

Ref.: Reid, 1965.

\**Stereum ochraceo-flavum* (Schw.) Sacc.

Syll. Fung. 6: 576, 1888.

BASIONYM: *Thelephora ochraceo-flava* Schw., 1832.

DISTRIBUTION: cosmopolitan.

Ref.: Lentz, 1955; Dueñas & Tellería, 1988; Hjortstam & Larsson, 1995; Eicker-Albert, 1998.

\**Stereum ostrea* (Blume & Nees) Fr.

Epicr. Syst. Mycol. p. 547. 1838.

BASIONYM: *Thelephora ostrea* Blume & Nees, 1826.

DISTRIBUTION: cosmopolitan.

Ref.: Lentz, 1955.




Figure 1 – Map of the State of Santa Catarina showing the areas of Atlantic Rain Forest and Parque Estadual da Serra do Tabuleiro.

### Acknowledgements

The authors thank Leif Ryvarden and Jorge E. Wright for reviewing the manuscript.

### Literature cited

- Azevedo CPL, Guerrero RT. 1993. Estudo biosistemático de espécies do gênero *Hymenochaete* (*Basidiomycetes*) no Rio Grande do Sul. *Insula* 22: 143-176.
- Batista MR. 2003. Caracterização geográfica. In: Fundação do Meio Ambiente. A natureza do Parque Estadual da Serra do Tabuleiro. Florianópolis: FATMA. 128 p.
- Blanchette RA. 1995. Degradation of the lignocellulose complex in wood. *Canadian Journal of Botany* 73 (suppl. 1): 999-1010.



- Burdsall HH, Miller OK. 1975. A reevaluation of *Panellus* and *Dictyopanus* (*Agaricales*). Nova Hedwigia Beiheft 51: 79-91.
- Cooke B. 1961. The genus *Schizophyllum*. Mycologia 53(6): 575-599.
- Dai YC. 2000. Changbai wood-rotting fungi 12. Species of *Hymenochaete* (*Basidiomycota*). Mycotaxon 76: 445-450.
- David A. 1982. Étude monographique du genre *Skeletocutis* (*Polyporaceae*). Naturaliste Can. (Rev. Écol. Syst.) 109: 235-272.
- Decock C, Figueroa SH, Ryvarden L. 2001. Studies in *Perenniporia contraria* and its presumed taxonomic synonym *Fomes subannosus*. Mycologia 93(1): 196-204.
- Dennis RWG. 1970. Fungus flora of Venezuela and adjacent countries. Kew Bulletin Additional Series III. 531 p.
- Dueñas M, Tellería MT. 1988. Catálogo de los corticiáceos y poliporáceos, S. L. (Aphyllophorales, Basidiomycotina), de la micoflora cántabro-astur. Madrid: Real Jardín Botánico, C. S. I. C. 262 p.
- Eicker-Albert LS. 1998. *Stereum* species (Stereaceae) of South Africa. South African Journal of Botany 64(1): 30-37.
- Eriksson J, Ryvarden L. 1976. The Corticiaceae of North Europe. Vol. 4 *Hyphodermella – Mycoacia*. Oslo, Fungiflora: 549-886.
- Fryar SC, Kirby GC, Hyde KD. 1999. Species abundance patterns of two wood decay basidiomycete communities. Fungal Diversity 3: 39-56.
- Furtado JS. 1981. Taxonomy of *Amauroderma* (Basidiomycetes, Polyporaceae). Memoirs of the New York Botanical Garden 34: 1-109.
- Gerber AL. 1996. Fungos xilófilos poróides (Aphyllophorales) no Morro da Lagoa da Conceição, Ilha de Santa Catarina, SC, Brasil. Insula 25: 3-68.
- Gerber AL, Loguercio-Leite C. 1997. New records of polypores (Aphyllophorales) from southern Brazil. Mycotaxon 62: 305-318.
- Gerber AL, Loguercio-Leite C. 2000. Polyporoid wood-rotting fungi (Basidiomycetes) II – New records from southern Brazil. Mycotaxon 76: 175-185.
- Gerber AL, Neves MA, Loguercio-Leite C. 1999. Some species of *Perenniporia* Murrill (*Poriales*, *Basidiomycotina*) from Southern Brazil. Revta. Brasil. Bot. 22(2): 185-193.
- Gilbert GS. 2002. Host specialization among wood-decay polypore fungi in a Caribbean mangrove forest. Biotropica 34(3): 396-404.
- Gilbert GS, Ferrer A, Carranza J. 2002. Polypore fungal diversity and host density in a moist tropical forest. Biodiversity and Conservation 11: 947-957.
- Gilbertson RL, Ryvarden L. 1986. North America polypores. *Abortiporus – Lindtneria*. Oslo, Fungiflora 1: 1-433.
- Gilbertson RL, Ryvarden L. 1987. North America polypores. *Megasporosporia – Wrightoporia*. Oslo, Fungiflora 2: 434-885.
- Gimms J. 1979. *Henningsia* (Polyporaceae) and a description of the type species. Mycologia 71(2): 305-309.
- Gimms J. 1980. The genus *Flaviporus* Murrill (Polyporaceae). Can. J. Bot. 58: 1578-1590.
- Gonçalves GVC, Loguercio-Leite C. 2001. Biodiversidade de fungos xilófilos (Basidiomycetes), na Unidade de Conservação Ambiental Desterro (UCAD), Ilha de Santa Catarina, SC, Brasil. Insula 30: 1-19.
- Gugliotta AM. 1997. *Polyporaceae* de mata ciliar da Estação Experimental e Reserva Biológica de Moji-Guaçu, SP, Brasil. Hoehnea 24(2): 89-106.
- Gulaid H, Ryvarden L. 1998. Two new species of *Amauroderma* (Ganodermataceae, Basidiomycetes). Mycologia Helvetica 10(1): 25-30.
- Guzmán G, Johnson D. 1974. Registros y especies nuevas de los hongos de Palenque, Chiapas. Bol. Soc. Mex. Mic. 8: 73-105.
- Hjortstam K, Larsson KH. 1995. Annotated check-list to genera and species of corticioid fungi (Aphyllophorales, Basidiomycotina) with special regards to tropical and subtropical areas. Windahlia 21: 1-75.

- Holmgren PK, Holmgren NH, Barnett LC. 1990. *Index Herbariorum*. Pt. 1. The Herbaria of the World. International Association for Plant Taxonomy, New York Botanical Garden, New York. 693 p.
- Hongo T. 1974. Agarics from Papua-New Guinea (2). Mycological reports from New Guinea and the Solomon Islands (27). Rept. Tottori Mycol. Inst. (Japan) 11: 29-41.
- Jejelowo OA, Abraham SP. 1998. A preliminary list of a larger fungi from kingwood forest Northeast of Houston, Texas. *Texas J. Sci.* 50(2): 143-148.
- Job DJ. 1985a. Basidiomicetos xilófilos de la región mesopotámica VI. Espécies del género *Hymenochaete* Lév. *Revista de Investigaciones Agropecuárias* 20(1): 143-165.
- Job DJ. 1985b. The South American collections of *Hymenochaete* Lév. (Aphyllphorales) in J. Rick's Herbarium. *Mycotaxon* 24: 227-235.
- Kirk PM, Cannon PF, David JC, Stalpers JA. 2001. *Dictionary of the Fungi*. 9ª edição; CAB International.
- Kotterman M, Heessels E, Jong E, Field JA. 1994. The physiology of anthracene biogradation by the white rot fungus *Bjerkandera* sp. Strain BOS55. *Applied Microbiology and Biotechnology* 42: 179-186.
- Larsen MJ, Cobb-Pouille LA. 1990. *Phellinus* (Hymenochaetaceae): a Survey of the World Taxa. Oslo, Fungiflora, Synopsis Fungorum 3, 206 p.
- Lentz PL. 1955. *Stereum* and allied genera of fungi in the Upper Mississippi Valley. Agriculture Monograph 24.
- Leonard AC. 1998. Two *Ganoderma* species compared. *Mycologist* 12(2): 65-68.
- Lodge DJ, Ammirati JF, O'Dell TE, Mueller GM. 2004. Collecting and describing macrofungi. In: Mueller GM, Bills GF, Foster MS. Biodiversity of fungi: inventory and monitoring methods. Elsevier Academic Press, Oxford, UK. 777 p.
- Loguercio-Leite C. 1991. Revisão histórica sobre fungos poliporóides (Aphyllphorales) xilófilos de Santa Catarina, Brasil. *Ínsula* 20: 3-10.
- Loguercio-Leite C. 1992. El género *Polyporus* (Polyporaceae) en la Isla de Santa Catarina, Santa Catarina, Brasil. *Bol. Soc. Argent. Bot.* 28(1-4): 27-36.
- Loguercio-Leite C. 1993. Polyporaceae II: *Trametes* Fr. Na Ilha de Santa Catarina, SC, Brasil. *Ínsula* 22: 3-20.
- Loguercio-Leite C. 1994. Polyporaceae na Ilha de Santa Catarina III: o genero *Hexagonia* Fr. *Ínsula* 23: 3-14.
- Loguercio-Leite C, Wright JE. 1991. New South American Pileate Polypores (Polyporales) from Santa Catarina Island, SC, Brazil. *Mycotaxon* 41(1): 167-172.
- Loguercio-Leite C, Wright JE. 1995. The Genus *Phellinus* (Hymenochaetaceae) on the Island of Santa Catarina, Brazil. *Mycotaxon* 54: 361-388.
- Loguercio-Leite C, Wright JE. 1998. *Diplomitoporus dilutabilis* a new species of *Polyporales* (Aphyllphorales) from Santa Catarina Island, Brazil. *Mycotaxon* 68: 47-51.
- Loguercio-Leite C, Gerber A, Ryvarden L. 1998. *Wrightoporia porilacerata*, a new species of pore fungi from southern Brazil. *Mycotaxon* 67: 251-255.
- Loguercio-Leite C, Gonçalves GVC, Ryvarden L. 2001. Studies in Neotropical polypores 13. *Ceriporiopsis cystidiata* sp. nov. *Mycotaxon* 79: 285-288.
- Loguercio-Leite C, Ryvarden L, Groppo C. 2002. Studies in neotropical polypores 16. *Rubroporus carneoporis* genus & species nova. *Mycotaxon* 83: 223-227.
- Ministério do Meio Ambiente. 2002. Biodiversidade Brasileira: avaliação e identificação de áreas e ações prioritárias para conservação, utilização sustentável e repartição de benefícios da biodiversidade brasileira. Brasília, MMA/SBF. 404 p.
- Mitra A. 1999. *Microporellus obovatus* in India and its cultural characteristics. *Mycotaxon* 70: 255-262.
- Morellato LPC, Haddad CFB. 2000. Introduction: the Brazilian Atlantic Forest. *Biotropica* 32(4b): 786-792.
- Núñez M, Ryvarden L. 2000a. East Asian Polypores. *Ganodermataceae* and *Hymenochaetaceae*. Oslo, Fungiflora, 168 p.

- Nuñez M, Ryvarden L. 2000b. East Asian Polypores. *Polyporaceae s. l.* Oslo, Fungiflora, 168 p.
- Pavlich M. 1976. *Ascomycetes y Basidiomycetes* del Peru. I Memórias del Museo de Historia Natural "Javier Prado" 17.
- Pegler DN. 1987. A revision of the *Agaricales* of Cuba 1. Species described by Berkeley & Curtis. Kew Bull. 42(3): 501-585.
- Rajchenberg M. 1987. Type studies of *Polyporaceae* (Aphyllphorales) described by J. Rick. Nord. J. Bot. 7: 533-568.
- Rayner ADM, Boddy L. 1988. Fungal decomposition of wood, its biology and ecology. John Wiley & Sons, Chichester. 587 p.
- Reid DA. 1965. A monograph of the stipitate stereoid fungi. Nova Hedwigia 18. 382 p.
- Rick J. 1938. Agarici Riograndenses. Lilloa II: 251-316.
- Rosário LA. 2003. Paisagem e Conservação. In: Fundação do Meio Ambiente. A natureza do Parque Estadual da Serra do Tabuleiro. Florianópolis: FATMA. 128 p.
- Rossmann AY, Tulloss RE, O'Dell TE, Thorn RG. 1998. Protocols for an all taxa biodiversity inventory of fungi in a Costa Rican conservation area. Parkway Publications, Inc., Boone, North Carolina. 163 p.
- Ryvarden L. 1972. Studies on the Aphyllphorales of the Canary Islands with a note on the genus *Perenniporia* Murr.. Norwegian Journal of Botany 19(2): 139-144.
- Ryvarden L. 1991. Genera of Polypores – Nomenclature and taxonomy. Synopsis Fungorum 5. Fungiflora, Oslo, 363 p.
- Ryvarden L, Johansen I. 1980. A preliminary polypore flora of East Africa. Oslo, Fungiflora, 630 p.
- Saber M. 1997. New records of *Agaricales* (Pleurotoid in habit) for Iran. Iranian Journal of Plant Pathology 33(3-4): 51-58.
- Setliff EC. 1972. The taxonomy and morphology of *Poria vincta*. Mycologia 64: 689-701.
- Silveira RMB, Guerrero RT. 1991. Aphyllphorales poliporóides (Basidiomycetes) do Parque Nacional de Aparados da Serra, Rio Grande do Sul. Boletim do Instituto de Biociências 48: 1-127.
- Singer R. 1975. The Agaricales in modern taxonomy. 2 ed. Rev. J. Cramer, Weinheim, 912 p.
- Soares SCS, Gugliotta AM. 1998. Criptógamos do Parque Estadual das Fontes do Ipiranga, São Paulo, SP. Fungos, 7: Aphyllphorales (Hymenochaetales). Hoehnea 25(1): 11-31.
- Smânia A, Marques CJS, EFA Smânia, Zanetti CR, Carobrez, SG, Tramonte R & Loguercio-Leite C. 2003. Toxicity and antiviral activity of cinnabarin obtained from *Pycnoporus sanguineus*. Phytol. Res. 17: 1069-1072.
- Steyaert RL. 1972. Species of *Ganoderma* and related genera mainly of the Bagor and Leiden Herbaria. Persoonia 7(1): 55-118.
- Steyaert RL. 1975. The concept and circumscription of *Ganoderma tornatum* (Pers.) Bres. Trans. Br. Mycol. Soc. 65(3): 451 – 567.
- Straatsma G, Krisai-Greilhuber I. 2003. Assemblage structure, species richness, abundance, and distribution of fungal fruit bodies in a seven year plot-based survey near Vienna. Mycological Research 107(5): 632-640.
- Tabanez AAJ, Viana VM. 2000. Patch structure within Brazilian Atlantic forest fragments and implications for conservation. Biotropica 32(4b): 925-933.
- Thomas WW. 1999. Conservation and monographic research on the flora of Tropical America. Biodiversity and Conservation 8: 1007-1015.
- Wagner T, Ryvarden L. 2002. Phylogeny and taxonomy of the genus *Phylloporia* (Hymenochaetales). Mycological Progress 1(1): 105-116.
- Zhang XQ. 1999. Fungal flora of tropical Guangxi, China: Aphyllphorales s.l. Mycotaxon 72: 371-376.