

Rabbit N. Bernick-

27.12.96

6/96

NT 528855

ENTERED

2.1

Viennotichia fumicola

Pilaira 11-9.5 x 7 anovata pale yellowish.

Schizotocia tetraspora 22.5 x 11.2.

13.1

Viem. fumicola

Sch. tetraspora

Ascochelus pale, st. vinaceous 17.5 x 11.2 ovoid,
(hr var overripe) Spores well ridged 19.3 x 11.2

Asci clavate 210 x 24

Paraphyses hyaline, st. clavate

14.1

up to 1.5 um diam

21 x 11

Spores basiconic. Asci broadly clavate 22.5 x 26.0 - 25.0

discharged spores 19.3-22.5 x 11-11.5

Cop. mixed appearing

Spores in ascos intensely purple. Ascos not noticeably blue in kg ascus contents purple/brown large unilateral mycelia.

probably large form of A. albidus

PTD

24.1

~~Hyphocya~~ up to 3m

to sign of furrowed web. Slight
virescent tinge to apical rim,
otherwise ± white.

30.1

Volvetella ciliata

Small whitestipitate Copriids, minutely hairy along
flattened apo 200 diam margin

Asci 8 spored biserial. Cylindrical clavate,
shrubly calved below 40-30-35 x 8

Spores ellipsoid. 4.5-5 x 3

to sign of any pore. Looks operulate.
no reaction with KI.

red oil Spores with faint
characteristics.

(Sent to B Spomer)

5.2.

Trichopanus cornus

from ~~8~~ below to 9-10. Pores 5-7 capitate,
Asci blue in KI

Spores are small size 17.7 x 11.2. gel sheath
all round.

Spores being ingested by amoebae

Ascidium - pale

22.5 x 11.2. unilatered gel

albicans

Dried 1.2.97