

30.3 T. hololeptcha hospita ✓ 25.7-29 x 9.6

31.3 Thelobolus stercomis Asus ✓ 280 x 190

5.4 Ascogonus hairs membrane below,
Same from lower part of apo
60, 12-13 at base, tapering to 6.5,
obuse apex smooth, 1-2 septate
Young, but base of globose cells 15-25
apo ca 300 diam.
Asci 112 x 32 spores not yet ripe.



5.4 "Sordaria polyspora" multispored, small-spored,
golden brown when young, dark grey/black ~~dark~~ when
ripe Asci ca 112 x 22 ✓
Pezizella immersed, tomentum dark grey, dark
valued stroma. Spores 4.5-5 x 3, 5:1
asymmetrical, P-T sloped ○

T. hospita ✓

Ascogonus Spores 8 x 3, none
ellipsoid than ~~Asus~~ form of Asus?
[might be for something else]
spores 12 x 6 broad ellipsoid / Asus form
some odd stone f-rod processes - can't

hairs at exactly where they come from

- seem to be at ascus tip after
spores liberated? brooked umbellate paraphysis apex?
Levellanus?

11.4

Sordaria 22.5 x 14.5 broad, papillate -
sheath not obvious humana

T. stercoraria Sch. vestitoria

26.4

Coprinus ✓ cardisporus Sch. vestitoria

Anoploperum ampullaceum very numerous ampullate hairs
all over to 65 long. Spores 32 x 16

T. bisporus ✓ Podospora ✓ appedrolata typical
perithecial hairs + slope

Conocochlea ✓ Satzigera 19.5 x 14.5 x 10 setae 48

1.5

P. appedrolata Sch. vestitoria

Sporewielder CAR 48-514 x 9.5-10
55- intermedia

Sporewielder CAR 35-42 x 7 antialis

5.5

Coprinus ✓ Cap. cardisporus Anop. ampullaceum

Sch. vestitoria P. appedrolata T. bisporus

Sp. antialis Con. Satzigera

Dried 5.5