

unidentified Lamington NP, 28°13'44"S; 153°7'28"E 26.10.99 75/99

small Qld  
mammal

inc 3.12

6.12 with spores Podospora broad clavate asc' ca 380 x 80-95  
broader in middle (?)

Spores, brown var-inflated, var-agg hairs  
Setae < 220  
128 x 256 spored.

Spores 12.5-13 x 8 Shell simple in 2nd perithecia  
Simple 2ys, var very apparent (apical gp)

Ascospora spores finely & minutely striate

only one ascus ca 125 x 14

spores 11-13 x 7 pores hyaline 3, set-cap.

examination of Polygord cells < 20 diam.

 agymnetic gel.

~~minutes~~ ✓

See 30.12

13.12. Polysporid Podospora asc' clavate ca 340 x 100  
280+ spores 11.5-12.5 x 8

Setae < 190

21.12 white furfuraceous Ascomata apothecia - immature.

large immersed perithecia Podospora polysporid

Setae all round, var particularly aggregated. Not inflated

Asci 260 x 85 broad saccate

Spores 11.2 x 8 tiny by app

ca 5 x 1 v. diff  
to see in active spores

**PACKET**



30.12 Podospora fuscescens hairy below, radially setae at  
 venter 480 diam semi immersed.  
 Saccate polysporous asci (256?) ca 350 x 95-100  
 Spores 12-13 x 8  
 14 4 x 2



Setae < 200  
 septate, minute + paler  
 at tip

Ascobolus pale green, furfuraceous scurfy, crenate  
 margin Asci ca 160 x 14-15-16

Spores finely & closely lined 12.5-13 x 6-6.5  
 very thin all round gel in II?

✓ crenulatus

7.1.00 Asc. crenulatus ✓

PACKET

Podospora ✓ radially setae polysporous setae  
 at venter non-aggregated, non-inflated. < 200  
 Spores small apical gp. 12.5-13 x 7-7.5



4-6 x 1-2

PACKET

13.1 Asc. crenulatus ✓

Podospora as above ✓

Dried 13.1

AB:- Ascobolus crenulatus ✓ [P] [P]  
Podospora sp. 1

+ Preussia furfurata  
 Coprinus luteus / baesporus