

16.8 Sporosmiella ♂ celled 40-48 x 6-6.5 ^{end cells rounded}
 St. longer than rest!
 height < 100 high dense black peridium
 Ascii ^{marked} ca 125 x 12-16 ^{tapers below to}
 irreg. biseriate, with one in base ^{long stalk.}
 but st. tapered. 3rd cell
 marginally shorter → under than the rest.
Oudemera ✓

20.8 Sp. immature 4 ER
 ca 34-35 ? check later

Conocheete Wimmer ✓ 10-11 x 8 x 5

24.8 Con Wimmer ✓ P. crustaceus ✓

29.8 Asc. condeleanii ♂ enveloping ? P. crustaceus ✓

Sp. 4 ER frequently 32-34 ✓ Mimra

Sp. Oudemera ✓ asci with long stalked
 tapers base 45-48

Also immature, with tapered asci/apical cells
 35 long. Lageniformis ?

31.8 Asciobolus Spores ± smooth Some reticulate, some with
 rare cracks, some about verrucae in places ✓

White apo. sp. 12.5-16 x 7.0-8 mostly condoleani
 Pattern, asynthetic gel Pores lichen
 3 im. v. 17. capitata or etc. Asc. 250 x 20
 excipular cells < 40 diam.

4.9 Asc. catenari / (dried)

Sp. minima ✓ R. crustaceus ✓

Coniochaeta S. sp 12 x 10 x 5 hymenium

10.9 Coprinus miser ✓ Coni hymenium
Setae 4/30

Sp. minima

15.9 Cop. miser ✓ Sp. minima ✓

① Coniochaeta hot very discard Spore 11.2 x 7 x 4.5
Setae ostiolar only → Asc ca 80 x 7 uniseriate

② Asper pentherm Spones 8-7 x 5 x 3
dense short hairs up to 16. at base

③ Asper 9.5 x 6.5 x 5 Asc ca 100 x 10 diam
spore short Setae cf. Lecanoplaea

Dried 15.9