

3.12 -

6.12

Pilaira Spore pale ellipsoid 9-13 x 6-8

10.12

Piplocephalis lessemiana X dry, brown  
dark - de - gout base fowl lead cell.

15.12

P. lessemiana ✓

Ascocarps woollyensis ✓ sm 55-65 x 19-22  
Sp. 12-13 x 4 setae < 50 long

KI - ve.

20.12

Cyp. strobliana ✓

A. woollyensis ✓

Volutella ciliata X sp. 5 x 2 setae thick-  
walled, Septate < 110 long.

25.12

Sch. tetrasporum 4-sp 20-21 x 12-12.5

Tel. ramos X

Asc. woollyensis -

Cyphella pink white 'firm' looking asc. 200-300 mic

8-sp. bisexate sp 11.5-12.5 x 7-8

Asc cylindrical 80-90 x 16.

Paras fine, not capitate, circinate ✓

Ascocarps variolos X

Asc 125-145 x 57-65

Spore wall 64-70 + 4c-50 spores rounded broad

sp. 12.5-13 x 5.5-6 - larger than other collections

Spores typically rounded - all relatives from central part

6.1 Cap sternover ✓ Sch. tetrasporum ✓

White apothecium ca 250 diam

Ascii ± cylindrical ca 60-70 x 8-9  
no neck or opercular spine  
spores biseriate long cylindrical, smooth,  
ca 12 x 2-3

?

10.1 Cap sternover ✓ Sch. tetrasporum ✓

Ascozome white, refined hair rough,  
or rot.

Smaller below Ascii with rounded apex,  
rot like A. wortmannii. Spore walls not fused

like A. w. Spores are ellipsoid, rather  
fusoid. Asc 90-100<sup>-105</sup> x 22-23-26 Spore with (16?) some  
with wall < 32 - spores. 10-11.5 x 6

exciator cells ± pyriform - sparse < 10 x 10  
no base of those cells.

appears 81. brownish ca 25 long.

could be Panspore or crovaiii ?

asci appear to be operculate - could refined  
hair be paraphyses - seem to have  
hyphae as well?

14.1 Cap sternover ✓

Ascozome? 32 - spored rounded spores

10-11 x 4.5-5

definitely operculate. 5

keve

pellet kept

Discarded 14.1