

Cattle. by river Hvita, Bryðjuholt, Fluðir, Hrunamannareppur, Iceland. 47/06
4359. W20.300; N64.189. 2.8.06. 64 m. Riverside pasture. Incubated 9.8.

- 11.8 ~~Pilobolus klenzi~~ ✓ sp. deep orange ellipsoid
13-15 x 9-10
- 15.8 Coprinus expanded to 18 m from stripe 8 cm long
○ 12.5-13 x 6.5-7 4-spored? ✓
- Sp. 4FR ✓ immature - not seen again - not recorded
- 17.8 Coprinus Scales - no sphaerocysts - No fructifera.
4-sp. no spores yet, just basidia
- 21.8 larval activity - sprayed
← spores 8 x 3.5
- Ascobolus imerius ✓ pellucidos ✓
- ~~Sch. zollneri~~ immature ♂ sp clear scales
- 25.8 Trichobolus microsporus ✓ Cop. pellucidos ✓
- Asc. imerius ✓ Cop. leptogonus ✓ scales + sphaerocysts
- Pezizopora decipiens ✓ ♂-sp 385-445 x 19-20
- Sch. canan ✓ ♂-sp food scales 24-25.5 x 14-14.5
- 1.9 T. microsporus ✓ Cop. lanabli ✓ P. decipiens ✓
- Lodoplasma canare ✓ sp 21-22.5 x 12-13 ellipsoid
mounting mixture - Asc: 270 x 50-52
- 320
- 2.9 Cop. lanabli 4-sp 10-11.5 x 6.5 rebridos ✓

22.8. Cop. nise, Cop. stellata clade.

24.9 Cop. nivar privata? T. microspora ✓

Cop. eddai no spheroid, no spores

~~Cop. nise~~ privata Sch. common ✓

25.9 Cop. nivar prairie-grass

Cop. nise/edda spores type-spores present on old fib

2.10 Cop. nivar ✓ T. microspora ✓ Sch. common ✓

Discarded 2.10